



## National CO<sub>2</sub> budgets (2015–2020) inferred from atmospheric CO<sub>2</sub> observations in support of the Global Stocktake

Brendan Byrne<sup>1</sup>, David F. Baker<sup>2</sup>, Sourish Basu<sup>3,4</sup>, Michael Bertolacci<sup>5</sup>, Kevin W. Bowman<sup>1,6</sup>, Dustin Carroll<sup>7,1</sup>, Abhishek Chatterjee<sup>1</sup>, Frédéric Chevallier<sup>8</sup>, Philippe Ciais<sup>8</sup>, Noel Cressie<sup>5,1</sup>, David Crisp<sup>1</sup>, Sean Crowell<sup>9</sup>, Feng Deng<sup>10</sup>, Zhu Deng<sup>11</sup>, Nicholas M. Deutscher<sup>12</sup>, Manvendra K. Dubey<sup>13</sup>, Sha Feng<sup>14</sup>, Omaira E. García<sup>15</sup>, David W. T. Griffith<sup>12</sup>, Benedikt Herkommer<sup>16</sup>, Lei Hu<sup>17,18</sup>, Andrew R. Jacobson<sup>17,18</sup>, Rajesh Janardanan<sup>19</sup>, Sujong Jeong<sup>20</sup>, Matthew S. Johnson<sup>21</sup>, Dylan B. A. Jones<sup>10</sup>, Rigel Kivi<sup>22</sup>, Junjie Liu<sup>1,23</sup>, Zhiqiang Liu<sup>24</sup>, Shamil Maksyutov<sup>19</sup>, John B. Miller<sup>17</sup>, Scot M. Miller<sup>25</sup>, Isamu Morino<sup>19</sup>, Justus Notholt<sup>26</sup>, Tomohiro Oda<sup>27,28</sup>, Christopher W. O'Dell<sup>2</sup>, Young-Suk Oh<sup>29</sup>, Hirofumi Ohyama<sup>19</sup>, Prabir K. Patra<sup>30</sup>, H el ene Peiro<sup>9</sup>, Christof Petri<sup>26</sup>, Sajeev Philip<sup>31</sup>, David F. Pollard<sup>32</sup>, Benjamin Poulter<sup>3</sup>, Marine Remaud<sup>8</sup>, Andrew Schuh<sup>2</sup>, Mahesh K. Sha<sup>33</sup>, Kei Shiomi<sup>34</sup>, Kimberly Strong<sup>10</sup>, Colm Sweeney<sup>17</sup>, Yao T e<sup>35</sup>, Hanqin Tian<sup>36,37</sup>, Voltaire A. Velazco<sup>12,38</sup>, Mihalis Vrekoussis<sup>39,26</sup>, Thorsten Warneke<sup>26</sup>, John R. Worden<sup>1</sup>, Debra Wunch<sup>10</sup>, Yuanzhi Yao<sup>36</sup>, Jeongmin Yun<sup>20</sup>, Andrew Zammit-Mangion<sup>5</sup>, and Ning Zeng<sup>28,4</sup>

<sup>1</sup>Jet Propulsion Laboratory, California Institute of Technology, Pasadena, CA, USA

<sup>2</sup>Cooperative Institute for Research in the Atmosphere, Colorado State University, Fort Collins, CO, USA

<sup>3</sup>NASA Goddard Space Flight Center, Global Modeling and Assimilation Office, Greenbelt, MD, USA

<sup>4</sup>Earth System Science Interdisciplinary Center, College Park, MD, USA

<sup>5</sup>School of Mathematics and Applied Statistics, University of Wollongong, Australia

<sup>6</sup>Joint Institute for Regional Earth System Science and Engineering, University of California, Los Angeles, CA, USA

<sup>7</sup>Moss Landing Marine Laboratories, San Jos e State University, Moss Landing, CA, USA

<sup>8</sup>Laboratoire des Sciences du Climat et de L'Environnement, LSCE/IPSL, CEA-CNRS-UVSQ, Universit e Paris-Saclay, 91191 Gif-sur-Yvette, France

<sup>9</sup>University of Oklahoma, Norman, OK, USA

<sup>10</sup>Department of Physics, University of Toronto, Toronto, Ontario, Canada

<sup>11</sup>Department of Earth System Science, Tsinghua University, Beijing, China

<sup>12</sup>Centre for Atmospheric Chemistry, School of Earth, Atmospheric and Life Sciences, University of Wollongong, Wollongong, NSW, Australia

<sup>13</sup>Earth System Observation, Los Alamos National Laboratory, Los Alamos, NM, USA

<sup>14</sup>Atmospheric Sciences and Global Change Division, Pacific Northwest National Laboratory, Richland, WA, USA

<sup>15</sup>Iza na Atmospheric Research Center (IARC), State Meteorological Agency of Spain (AEMet), Tenerife, Spain

<sup>16</sup>Institut for Meteorology and Climate Research (IMK-ASF), Karlsruhe Institute of Technology (KIT), Karlsruhe, Germany

<sup>17</sup>NOAA Global Monitoring Laboratory, Boulder, CO, USA

<sup>18</sup>Cooperative Institute for Research in Environmental Sciences, University of Colorado Boulder, Boulder, CO, USA

<sup>19</sup>Satellite Observation Center, Earth System Division, National Institute for Environmental Studies, Tsukuba, Japan

<sup>20</sup>Department of Environmental Planning, Graduate School of Environmental Studies, Seoul National University, Seoul, Republic of Korea

<sup>21</sup>NASA Ames Research Center, Moffett Field, CA, USA

<sup>22</sup>Space and Earth Observation Centre, Finnish Meteorological Institute, Sodankyl a, Finland

<sup>23</sup>Division of Geological and Planetary Sciences, California Institute of Technology, Pasadena, CA, USA

<sup>24</sup>Laboratory of Numerical Modeling for Atmospheric Sciences & Geophysical Fluid Dynamics, Institute of Atmospheric Physics, Chinese Academy of Sciences, Beijing, China



- <sup>25</sup>Department of Environmental Health and Engineering, Johns Hopkins University, Baltimore, MD 21218, United States of America
- <sup>26</sup>Institute of Environmental Physics, University of Bremen, Bremen, Germany
- <sup>27</sup>Earth from Space Institute, Universities Space Research Association, Columbia, MD, USA
- <sup>28</sup>Department of Atmospheric and Oceanic Science, University of Maryland, USA
- <sup>29</sup>Global Atmosphere Watch Team, Climate Research Department, National Institute of Meteorological Sciences, Republic of Korea
- <sup>30</sup>Research Institute for Global Change, Japan Agency for Marine-Earth Science and Technology (JAMSTEC), Yokohama, 236-0001, Japan
- <sup>31</sup>Centre for Atmospheric Sciences, Indian Institute of Technology Delhi, New Delhi, India
- <sup>32</sup>National Institute of Water & Atmospheric Research Ltd (NIWA), Lauder, New Zealand
- <sup>33</sup>Royal Belgian Institute for Space Aeronomy (BIRA-IASB), Brussels, Belgium
- <sup>34</sup>Japan Aerospace Exploration Agency (JAXA), Tsukuba, Japan
- <sup>35</sup>Laboratoire d'Etudes du Rayonnement et de la Matière en Astrophysique et Atmosphères (LERMA-IPSL), Sorbonne Université, CNRS, Observatoire de Paris, PSL Université, 75005 Paris, France.
- <sup>36</sup>International Center for Climate and Global Change Research, College of Forestry, Wildlife and Environment, Auburn University, Auburn, AL 36849, USA
- <sup>37</sup>Schiller Institute for Integrated Science and Society, and Department of Earth and Environmental Sciences, Boston College, Chestnut Hill, MA 02467, USA
- <sup>38</sup>Deutscher Wetterdienst (DWD), Hohenpeissenberg, Germany.
- <sup>39</sup>Climate and Atmosphere Research Center (CARE-C), The Cyprus Institute, Nicosia, Cyprus

**Correspondence:** Brendan Byrne (brendan.k.byrne@jpl.nasa.gov)

**Abstract.** Accurate accounting of emissions and removals of CO<sub>2</sub> is critical for the planning and verification of emission reduction targets in support of the Paris Agreement. Here, we present a pilot dataset of country-specific net carbon exchange (NCE; fossil plus terrestrial ecosystem fluxes) and terrestrial carbon stock changes aimed at informing countries' carbon budgets. These estimates are based on “top-down” NCE outputs from the v10 Orbiting Carbon Observatory (OCO-2) modeling inter-comparison project (MIP), wherein an ensemble of inverse modeling groups conducted standardized experiments assimilating OCO-2 column-averaged dry-air mole fraction (X<sub>CO<sub>2</sub></sub>) retrievals (ACOS v10), in situ CO<sub>2</sub> measurements, or combinations of these data. The v10 OCO-2 MIP NCE estimates are combined with “bottom-up” estimates of fossil fuel emissions and lateral carbon fluxes to estimate changes in terrestrial carbon stocks, which are impacted by anthropogenic and natural drivers. These flux and stock change estimates are reported annually (2015–2020) as both a global 1° × 1° gridded dataset and as a country-level dataset. Across the v10 OCO-2 MIP experiments, we obtain increases in the ensemble median terrestrial carbon stocks of 3.29–4.58 PgCO<sub>2</sub> yr<sup>-1</sup> (0.90–1.25 PgC yr<sup>-1</sup>). This is a result of broad increases in terrestrial carbon stocks across the northern extratropics, while the tropics generally have stock losses but with considerable regional variability and differences between v10 OCO-2 MIP experiments. We discuss the state of the science for tracking emissions and removals using top-down methods, including current limitations and future developments towards top-down monitoring and verification systems.

15 *Copyright statement.* ©2022. California Institute of Technology, government sponsorship acknowledged.



## 1 Introduction

To reduce the risks and impacts of climate change, the Paris Agreement aims to limit the global average temperature increase to well below 2 °C above pre-industrial levels and pursue efforts to limit these increases to less than 1.5 °C. To this end, each Party to the Paris Agreement agreed to prepare and communicate successive Nationally Determined Contributions (NDCs) of greenhouse gas (GHG) emission reductions. Collective progress toward this goal of the Paris Agreement is evaluated in Global Stocktakes (GSTs), which are conducted at five-year intervals; the first GST is scheduled in 2023. The outcome of each GST is then used as input, or as a “ratchet mechanism”, for new NDCs that are meant to encourage greater ambition.

In support of the first GST, Parties to the Paris Agreement are compiling inventories of GHG emissions and removals to inform their progress toward the emission-reduction targets in their individual NDCs. These inventories are generally estimated using “bottom-up” approaches, wherein CO<sub>2</sub> emission estimates are based on activity data and emission factors while CO<sub>2</sub> removals by sinks are based on inventories of carbon stock changes and models. This approach allows for explicit characterization of CO<sub>2</sub> emissions and removals into the five main sectors specified in the 2006 IPCC Guidelines for National Greenhouse Gas Inventories (IPCC, 2006): Energy, Industrial Processes and Product Use (IPPU), Agriculture, Forestry and Other Land Use (AFOLU), Waste, and Other. Bottom-up methods can provide precise and accurate country-level emission estimates when the activity data and emission factors are well quantified and understood, such as for the fossil fuel combustion category of the energy sector in many countries. However, these estimates can also have considerable uncertainty when the emission processes are challenging to quantify (such as for AFOLU) or if the activity data are inaccurate or missing. In addition, these estimates do not capture carbon emissions and removals from unmanaged systems, which are not directly considered in the Paris Agreement, but nevertheless impact the global carbon budget and growth rate of atmospheric CO<sub>2</sub>.

As a complement to these accounting-based inventory efforts, an independent “top-down” assessment of net surface-atmosphere CO<sub>2</sub> fluxes may be obtained from ground-based, airborne and space-based observations of atmospheric CO<sub>2</sub> mole fractions. These top-down methods have undergone rapid improvements in recent years, as recognized in the 2019 Refinement to the 2006 IPCC Guidelines for National Greenhouse Gas Inventories (IPCC, 2019). And, although these methods were not deemed to be a standard tool for verification of conventional inventories, a number of countries (UK, Switzerland, USA, and New Zealand) have adopted atmospheric inverse modeling as a verification system in national inventory reports. Initially, these countries have focused on non-CO<sub>2</sub> gasses (e.g., EPA, 2022), but top-down assessments of the CO<sub>2</sub> budget are now underdevelopment in New Zealand (<https://niwa.co.nz/climate/research-projects/carbon-watch-nz>). Furthermore, significant investments towards building anthropogenic CO<sub>2</sub>-emissions monitoring and verification support capacity are ongoing within the European Commission’s Copernicus Program (see Sect. 8.2.1).

In top-down CO<sub>2</sub> flux estimation, the net surface-atmosphere CO<sub>2</sub> fluxes are inferred from atmospheric CO<sub>2</sub> observations using state-of-the-art atmospheric CO<sub>2</sub> inversion systems (e.g., Peiro et al., 2022). This approach provides spatially- and temporally-resolved estimates of surface-atmosphere fluxes for land and ocean regions from which country-level annual land-atmosphere CO<sub>2</sub> fluxes can be estimated. The impact of fossil fuel (and usually fire CO<sub>2</sub> emissions) on the observations are accounted for in the inversions by prescribing maps of those emissions and assuming that they are perfectly known. Thus,



50 fossil fuel and fire CO<sub>2</sub> emissions are not diagnosed yet by these inversions, but net surface–atmosphere CO<sub>2</sub> fluxes from the terrestrial biosphere and oceans are. Terrestrial carbon stock changes can then be calculated by combining net surface–atmosphere CO<sub>2</sub> fluxes with estimates of fossil fuel emissions and horizontal (“lateral”) fluxes occurring within the terrestrial biosphere or between the land and ocean (Kondo et al., 2020). One example of a lateral flux is harvested agricultural products, where carbon is sequestered from the atmosphere by photosynthesis in one region but then this carbon is harvested and exported  
55 to another region as agricultural products. Similarly, carbon sequestered by photosynthesis in a forest can be leached away by streams and rivers, and then exported to the ocean. These lateral carbon fluxes are not directly identifiable in atmospheric CO<sub>2</sub> measurements, but accounting for their impact is required in order to convert net land fluxes into stock changes. These estimated terrestrial carbon stock changes reflect the combined impact of direct anthropogenic activities and changes to both managed and unmanaged ecosystems in response to rising CO<sub>2</sub>, climate change, and disturbance events (such as fires).

60 The top-down budgets presented here extend several previous studies that have developed approaches to compare inversion results to United Nation Framework Convention of Climate Change (UNFCCC) inventories. Ciais et al. (2021) proposed a protocol for reporting bottom-up and top-down fluxes so that they can be compared consistently. Chevallier (2021) noted that inversion results for terrestrial CO<sub>2</sub> fluxes should be restricted to managed lands and applied a managed land mask to the gridded fluxes of the CAMS CO<sub>2</sub> inversions for the comparison to UNFCCC values in ten large countries or groups of  
65 countries. Deng et al. (2022) compared CO<sub>2</sub>, CH<sub>4</sub> and N<sub>2</sub>O fluxes from inversion ensembles available from the Global Carbon Project. For CO<sub>2</sub>, they used six CO<sub>2</sub> flux estimates from inverse models that assimilated measurements from the global air-sample network, filtered their results over managed lands and corrected them for CO<sub>2</sub> fluxes induced by lateral processes to compare with carbon stock changes reported to the UNFCCC by a set of 12 countries. We expand upon these previous studies by providing top-down CO<sub>2</sub> budgets from the v10 Orbiting Carbon Observatory Model Intercomparison Project (v10 OCO-2  
70 MIP), wherein an ensemble of inverse modeling groups conducted standardized experiments assimilating OCO-2 column-averaged dry-air mole fraction ( $X_{\text{CO}_2}$ ) retrievals (retrieved with version 10 of the Atmospheric CO<sub>2</sub> Observations from Space (ACOS) full-physics retrieval algorithm), in situ CO<sub>2</sub> measurements, or combinations of these data. This allows us to quantify the sensitivity of top-down carbon budget estimates to the inversion modeling system and the atmospheric CO<sub>2</sub> dataset used to constrain flux estimates.

75 This paper is outlined as follows. The remainder of Sect. 1 describes the objectives of this work (Sect. 1.1) and provides background information on both the global carbon cycle (Sect. 1.2) and top-down atmospheric CO<sub>2</sub> inversions (Sect. 1.3). Section 2 defines the carbon cycle fluxes of interest. Section 3 describes the flux datasets and their uncertainties, including: fossil fuel emissions, the v10 OCO-2 MIP, riverine fluxes, wood fluxes, crop fluxes, and the net terrestrial carbon stock loss. Section 4 provides an evaluation of the v10 OCO-2 MIP flux estimates. Section 5 presents two metrics for interpreting the  
80 top-down constraints on the CO<sub>2</sub> budget. Section 6 gives a description of the dataset, Sect. 7 shows the characteristics of the dataset, and Sect. 8 discusses current limitations and future directions. Finally, Sect. 9 gives the conclusions of this study.



## 1.1 Objectives

This is a pilot project designed to start a dialogue between the top-down research community, inventory compilers, and the GHG assessment community to identify ways that top-down CO<sub>2</sub> flux estimates can help inform country-level carbon budgets (see Worden et al. (2022) for a similar pilot methane dataset). To meet this objective, the primary goal of this work is to provide two products: (1) annual net surface–atmosphere CO<sub>2</sub> fluxes and (2) annual changes in terrestrial carbon stocks. These products are provided annually over the six-year period 2015–2020 on both a 1° × 1° global grid and as country-level totals with error characterization.

These products are intended to be used to help inform inventory development and identify areas for future research in both top-down and bottom-up approaches. Including, informing strategies for operational top-down carbon cycle products that can be used for tracking combined changes in managed and unmanaged carbon stocks and help quantify the impact of emission reduction activities.

## 1.2 Overview of the carbon cycle

The carbon cycle describes the movement of carbon between various reservoirs in the Earth system. Carbon cycles between the atmosphere, oceans, and terrestrial biosphere through a number of pathways, while the burning of fossil fuels and cement production release geologic carbon to the atmosphere ( $40.0 \pm 3.3 \text{ PgCO}_2 \text{ yr}^{-1}$  or  $10.9 \pm 0.9 \text{ PgC yr}^{-1}$  over 2010–2019; Canadell et al., 2021). On an annual net basis, roughly half of the emitted CO<sub>2</sub> from anthropogenic sources is absorbed by terrestrial ecosystems and oceans (Friedlingstein et al., 2022), reducing the rate of atmospheric CO<sub>2</sub> increase ( $18.7 \pm 0.08 \text{ PgCO}_2 \text{ yr}^{-1}$  or  $5.1 \pm 0.02 \text{ PgC yr}^{-1}$  over 2010–2019; Canadell et al., 2021). Here we briefly review the movement of carbon between the reservoirs, and how these processes are modulated by human activities.

Fluxes of carbon between the atmosphere and ocean are driven by the difference in partial pressures of CO<sub>2</sub> between seawater and air, resulting in roughly balancing fluxes from the ocean-to-atmosphere and atmosphere-to-ocean of  $\sim 293 \text{ PgCO}_2 \text{ yr}^{-1}$  ( $\sim 80 \text{ PgC yr}^{-1}$ ) each way (Ciais et al., 2013), with a residual net atmosphere-to-ocean flux due to increasing atmospheric CO<sub>2</sub> ( $9.2 \pm 2.2 \text{ PgCO}_2 \text{ yr}^{-1}$  or  $2.5 \pm 0.6 \text{ PgC yr}^{-1}$  over 2010–2019; Canadell et al., 2021). Regional variations in the solubility and saturation of CO<sub>2</sub> in ocean waters drive net fluxes, with net fluxes to the atmosphere in upwelling regions, such as the eastern boundary of basins and in equatorial zones (McKinley et al., 2017). Meanwhile, there is net removals by the ocean in western boundary currents and at extratropical latitudes (McKinley et al., 2017). Within the oceans, circulation patterns, mixing, and biologic activity act to redistribute carbon.

On land, terrestrial ecosystems remove atmospheric carbon through photosynthesis, referred to as Gross Primary Production (GPP) (Fig. 1). GPP draws roughly  $440 \text{ PgCO}_2 \text{ yr}^{-1}$  ( $120 \text{ PgC yr}^{-1}$ ) from the atmosphere (Anav et al., 2015). Roughly half of this carbon is emitted back to the atmosphere by plants through autotrophic respiration, while the remaining carbon is used to generate plant biomass and is referred to as Net Primary Production (NPP). On an annual basis, the carbon sequestered through NPP is roughly balanced by carbon loss through a number of processes. The largest of these processes is heterotrophic respiration, which is the respiratory emission of CO<sub>2</sub> (from the dead organic matter and soil carbon pools) by heterotrophic




115 organisms, and accounts for 82–95% of NPP (Randerson et al., 2002). The combination of heterotrophic and autotrophic  
respiration is called ecosystem respiration ( $R_{eco}$ ). The remaining processes have smaller magnitudes, but are still critical for  
determining the carbon balance of ecosystems. Biomass burning, the emission of carbon to the atmosphere through combustion,  
releases roughly  $7.3 \text{ PgCO}_2 \text{ yr}^{-1}$  ( $2 \text{ PgC yr}^{-1}$ ) to the atmosphere on an annual basis, but with considerable interannual  
120 of carbon monoxide (CO), methane ( $\text{CH}_4$ ) and other biologic volatile organic compounds (BVOCs), which are oxidized to  $\text{CO}_2$   
in the atmosphere. Rivers move carbon in the form of dissolved inorganic carbon (DIC), dissolved organic carbon (DOC), and  
particulate organic carbon (POC). This carbon of terrestrial origin is partly transported to the open ocean, partly released to the  
atmosphere from inland waters and estuaries, and partly buried in aquatic or marine sediments. Finally, anthropogenic activities  
such as harvesting of crop and wood products result in lateral transport of carbon, such that the removal of atmospheric  $\text{CO}_2$   
125 through NPP and emission of atmospheric  $\text{CO}_2$  through respiration (e.g., decomposition in a landfill) or combustion (e.g.,  
burning of biofuels) occurs in different regions. See Fig. 1 for an illustration of these fluxes.

Globally, there is a long-term net uptake of atmospheric  $\text{CO}_2$  by the land (approximately  $-6.6 \text{ PgCO}_2 \text{ yr}^{-1}$  or  $-1.8 \text{ PgC}_2 \text{ yr}^{-1}$   
over 2010–2019; Canadell et al., 2021), which is the residual of an emission due to net land use change ( $5.9 \pm 2.6 \text{ PgCO}_2 \text{ yr}^{-1}$   
or  $1.6 \pm 0.7 \text{ PgC yr}^{-1}$  over 2010–2019; Canadell et al., 2021) and removal by other terrestrial ecosystems ( $12.6 \pm 3.3 \text{ PgCO}_2 \text{ yr}^{-1}$   
130 or  $3.4 \pm 0.9 \text{ PgC yr}^{-1}$  over 2010–2019; Canadell et al., 2021). This removal is partially driven by direct feedbacks between  
increasing  $\text{CO}_2$  and the biosphere, such as  $\text{CO}_2$  fertilization of photosynthesis and increased water use efficiency. Carbon-  
climate feedbacks also lead to both increases and decreases in terrestrial carbon stocks: for example, warming at high latitudes  
leads to a more productive biosphere but it also leads to increased plant and soil respiration (Kaushik et al., 2020; Walker et al.,  
2021; Canadell et al., 2021; Crisp et al., 2022). In addition, the release of nitrogen through anthropogenic energy and fertilizer  
135 use may also drive increased carbon sequestration by the terrestrial biosphere. Regrowth of forests in previously cleared areas,  
especially in the extratropics, is also thought to be an important uptake term (Kondo et al., 2018; Cook-Patton et al., 2020).  
Currently, the relative impact of each of these contributions to long-term terrestrial carbon sequestration is poorly known, and  
likely varies between biomes and climates.

While the existence of a long-term global land sink is supported through a number of lines of evidence (Ballantyne et al.,  
140 2012; Keeling and Graven, 2021), regional-scale emissions and removals are less well quantified. Regional-scale carbon se-  
questration can differ substantially from the global mean and can be impacted by the regional climate, disturbance events  
(Frank et al., 2015; Wang et al., 2021), and anthropogenic activities (Caspersen et al., 2000; Harris et al., 2012). The need to  
better quantify regional-scale emissions and removals of carbon has motivated much of the recent expansion of in situ  $\text{CO}_2$   
observing networks, the launch of space-based  $\text{CO}_2$  observing systems, and the development of  $\text{CO}_2$  inversion systems.

### 145 1.3 Background on atmospheric $\text{CO}_2$ inversions

Atmospheric  $\text{CO}_2$  inversions estimate the underlying net surface–atmosphere  $\text{CO}_2$  fluxes from atmospheric  $\text{CO}_2$  observations,  
and this is what is meant by the “top-down” approach (Bolin and Keeling, 1963; Tans et al., 1990; Enting et al., 1995; Gurney  
et al., 2002; Peiro et al., 2022). In this approach, an atmospheric chemical transport model (CTM) is employed to relate



**Figure 1.** CO<sub>2</sub> is removed from the atmosphere through photosynthesis (GPP) and then emitted back to the atmosphere through a number of processes. Three processes move carbon laterally on Earth’s surface, such that emissions of CO<sub>2</sub> occur in a different region than removals: (1) Agriculture; harvested crops are transported to urban areas and to livestock, which are themselves exported to urban areas. CO<sub>2</sub> is respired to the atmosphere in livestock or urban areas. (2) Forestry; logged carbon is transported to urban and industrial areas, then emitted to through decomposition in a landfill or combustion as a biofuel. (3) Water cycle; carbon is leached from soils into water bodies, such as lakes. The carbon is then either deposited, released to the atmosphere, or transported to the ocean (Regnier et al., 2022). Arrows show carbon fluxes and colors indicate whether the flux is associated with (grey) fossil fuel emissions, (dark green) ecosystem metabolism, (red) biomass burning, (light green) forestry, (yellow) agriculture, or (blue) the water cycle. Semi-transparent arrows show fluxes that move between the surface and atmosphere, while solid arrows show fluxes that move between land regions. Dashed arrows show surface–atmosphere fluxes of reduced carbon species that are oxidized to CO<sub>2</sub> in the atmosphere. For simplicity, a cement carbonation sink, volcano emissions, and a weathering sink are not included in this figure.

150 surface–atmosphere CO<sub>2</sub> fluxes to observed atmospheric CO<sub>2</sub> mole fractions. As an inverse problem, the upwind CO<sub>2</sub> fluxes are estimated from the downwind observed CO<sub>2</sub> mole fractions. The surface CO<sub>2</sub> fluxes are adjusted so that forward-simulated CO<sub>2</sub> mole fractions better match the CO<sub>2</sub> measurements while considering the uncertainty statistics on the observations, transport, and prior surface fluxes.



The atmospheric CO<sub>2</sub> inversion problem is generally ill-posed, such that the solution is underdetermined by the observational constraints. In this case, additional information is required to produce a unique solution and prevent overfitting of the data (Lawson and Hanson, 1974; Tarantola, 2005). Typically, this is performed using Bayesian inference, where prior mean fluxes and their uncertainties provide additional information required to estimate fluxes (Rayner et al., 2019). Prior mean fluxes of net ecosystem exchange are usually obtained from dynamic global vegetation models, while prior mean air-sea fluxes are derived from surface water partial pressure of CO<sub>2</sub> (pCO<sub>2</sub>) datasets or from ocean models (e.g., Peiro et al., 2022). The resulting posterior flux estimates combine the constraints on surface fluxes from atmospheric CO<sub>2</sub> data with the prior knowledge of the fluxes. If there is a high density of assimilated CO<sub>2</sub> observations, then the posterior fluxes will be more strongly impacted by the assimilated data, whereas, in regions with sparse observational coverage, the posterior fluxes will generally remain similar to the prior fluxes (assuming similar prior flux uncertainties across regions).

Measurements of atmospheric CO<sub>2</sub> best inform diffuse biosphere–atmosphere fluxes on large spatial scales. This is because CO<sub>2</sub> has a long atmospheric lifetime, such that the perturbation to atmospheric CO<sub>2</sub> due to emissions and removals from individual processes and locations gets mixed in the atmosphere (Gloor et al., 2001; Liu et al., 2015). For example, the measurements of CO<sub>2</sub> at Mauna Loa, Hawaii, provide a good estimate of the global-scale changes of CO<sub>2</sub> surface fluxes. Inferring smaller-scale flux signals requires a high density of CO<sub>2</sub> observations (to capture gradients in atmospheric CO<sub>2</sub>) and accurate modeling of atmospheric transport (to relate the measurements with surface fluxes). The accuracy of flux estimates depend on a number of factors, particularly the accuracy and precision of the data, transport model, and prior constraints. Stringent requirements on the accuracy of space-based column-averaged dry-air mole fraction ( $X_{\text{CO}_2}$ ) retrievals are required to infer surface fluxes (Chevallier et al., 2005a; Miller et al., 2007). Biases in  $X_{\text{CO}_2}$  retrievals from the Orbiting Carbon Observatory (OCO-2) related to spectroscopic errors, solar zenith angle, surface properties, and atmospheric scattering by clouds and aerosols have been identified (Wunch et al., 2017b). However, intensive research has reduced retrieval errors over time (O’Dell et al., 2018; Kiel et al., 2019). As will be shown in Sect. 4.1, biases in OCO-2  $X_{\text{CO}_2}$  retrievals over land are thought to be relatively small, although regionally structured biases may be present. However, OCO-2  $X_{\text{CO}_2}$  retrievals over oceans may contain more large-scale spatially coherent retrieval errors that can adversely impact flux estimates.

Accurate atmospheric transport is critical for correctly relating surface–atmosphere fluxes to observations. Due to computational constraints, CTMs are typically run offline with coarsened meteorological fields relative to the parent Numerical Weather Prediction model, which has been shown to introduce systematic transport errors in some configurations (Yu et al., 2018; Stanevich et al., 2020). In addition, these offline CTMs have been shown to have large-scale systematic differences in transport associated with the implementation of transport algorithms (Schuh et al., 2019, 2022). These errors appear to be of the same order as the retrieval biases, although the patterns in time and space are different. Systematic errors related to model transport (and errors in prior information) can partially be accounted for by performing multiple inversions that differ in CTM and prior constraints employed. This motivates inversion model intercomparison projects (MIPs), such as the OCO-2 MIP project (see Sect. 3.2; Crowell et al., 2019; Peiro et al., 2022). From these ensembles of inversions, estimates of both systematic errors (accuracy) and random errors (precision) can be obtained from the model spread.





## 2 Definitions

In this work, we focus on the carbon budget of Earth's land area, including aquatic systems such as rivers and lakes. In particular, we consider fluxes of carbon between the land and the atmosphere, and lateral carbon transport processes on land and between the land and ocean. We define the following annual net carbon fluxes (see Fig. 2 for an schematic representation of these fluxes):

- **Fossil fuel and cement emissions (FF)**: The burning of fossil fuels and release of carbon due to cement production, representing a flux of carbon from the land surface (geologic reservoir) to the atmosphere.
- **Net Biosphere exchange (NBE)**: Net flux of carbon from the terrestrial biosphere to the atmosphere due to biomass burning (BB) and ecosystem respiration ( $R_{\text{eco}}$ ) minus Gross Primary Production (GPP) (i.e.,  $\text{NBE} = \text{BB} + R_{\text{eco}} - \text{GPP}$ ). It includes both anthropogenic processes (e.g., deforestation, reforestation, farming) and natural processes (e.g., climate-variability-induced carbon fluxes, disturbances, recovery from disturbances).
- **Terrestrial Net Carbon Exchange (NCE)**: Net flux of carbon from the surface to the atmosphere. For land, NCE can be defined as:


$$\text{NCE} = \text{NBE} + \text{FF} \quad (1)$$

- **Lateral crop flux ( $F_{\text{crop trade}}$ )**: The lateral flux of carbon in (positive) or out (negative) of a region due to agriculture.
- **Lateral wood flux ( $F_{\text{wood trade}}$ )**: The lateral flux of carbon in (positive) or out (negative) of a region due to wood product harvesting and usage.
- **Lateral river flux ( $F_{\text{rivers export}}$ )**: The lateral flux of carbon in (positive) or out (negative) of a region transported by the water cycle.
- **Net terrestrial carbon stock loss ( $\Delta C_{\text{loss}}$ )**: Positive values indicate a loss (decrease) of terrestrial carbon stocks (organic matter stored on land), including above- and below-ground biomass in ecosystems and biomass contained in anthropogenic products (lumber, cattle, etc). This is calculated as:

$$\Delta C_{\text{loss}} = \text{NBE} - F_{\text{crop trade}} - F_{\text{wood trade}} - F_{\text{rivers export}} \quad (2)$$

- **Net terrestrial carbon stock gain ( $\Delta C_{\text{gain}}$ )**: Positive values indicate a gain (increase) of terrestrial carbon stocks, and is the negative of  $\Delta C_{\text{loss}}$ :

$$\Delta C_{\text{gain}} = -\Delta C_{\text{loss}} \quad (3)$$



**Figure 2.** Carbon fluxes for a given land region, such as a country. Boxes with solid backgrounds show reservoirs of carbon. Arrows with hatched shading show fluxes between reservoirs. NCE is underlined to emphasize that this quantity is estimated from the atmospheric CO<sub>2</sub> measurements using top-down methods. Italicized quantities are obtained from bottom-up datasets ( $FF$ ,  $F_{crop\ trade}$ ,  $F_{wood\ trade}$ ,  $F_{rivers\ export}$ ). Bold quantities are derived in this study from the top-down and bottom-up datasets ( $NBE$ ,  $\Delta C_{gain}$ ,  $\Delta C_{loss}$ ).

## 2.1 Country and regional aggregation

To aggregate gridded  $1^\circ \times 1^\circ$  flux estimates to country totals we use a country mask (Center for International Earth Science  
 215 Information Network - CIESIN - Columbia University, 2018). We also provide  $NCE$  and  $\Delta C_{loss}$  estimates for several country groupings. A number of regional intergovernmental organizations are included: the Association of Southeast Asian Nations (ASEAN), the African Union (AU) and each of its sub-regions (North, South, West, East, and Central), the Community of Latin American and Caribbean States plus Brazil (CELAC+Brazil), the Economic Cooperation Organization (ECO), the European Union (EU), and the South Asian Association for Regional Cooperation (SAARC). We also include some geographic regions,



220 specifically North America, the Middle East and Europe. Countries included in these groupings are listed in the supplementary materials (Text S1).

### 3 Flux datasets

Here, we describe the methodologies and datasets for estimating FF (Sect. 3.1), NCE (Sect. 3.2), country-level  $F_{\text{rivers export}}$  (Sect. 3.3), country-level  $F_{\text{crop trade}}$  and country-level  $F_{\text{wood trade}}$  (Sect. 3.4). Gridded lateral fluxes are estimated using a  
225 somewhat different approach, and are described in Sect. 3.5. Finally, we describe how these data are used to estimate  $\Delta C_{\text{loss}}$  (Sect. 3.6).

#### 3.1 Fossil fuel and cement emissions

Gridded  $1^\circ \times 1^\circ$  fossil  $\text{CO}_2$  emissions, including from cement production, are calculated as follows. Monthly gridded emissions up to 2019 are taken from the 2020 version of the Open-source Data Inventory for Anthropogenic  $\text{CO}_2$  (ODIAC2020, 2000–  
230 2019) emission data product (Oda and Maksyutov, 2011; Oda et al., 2018). The 2020 emissions were not part of ODIAC, but were projected using the Carbon Monitor (CM) emission data product (<https://carbonmonitor.org/>, downloaded 19th May 2021). For each month in 2020 and later, the ratio between that month's emissions and the emissions from the same month in 2019 was calculated from the CM emission data. Since CM provides daily emissions per sector for a handful of major emitting countries and the globe, CM emissions are summed over sectors and days in each month to create monthly total emissions per  
235 named country and the rest of the world (RoW). The ratio of each (post-2019) month's emission to the same month in 2019 is then calculated per named country and RoW, then distributed over a  $1^\circ \times 1^\circ$  grid assuming homogeneity of the ratio over each named country and RoW. 2019 ODIAC emissions for that month are then multiplied by the ratio to generate  $1^\circ \times 1^\circ$  monthly emissions after 2019. While this method loses the information of day-to-day variability provided by CM, this is a conscious choice to be consistent over the entire inversion period. Finally, we impose day-of-week and hour-of-day variations on these  
240 fluxes following the Temporal Improvements for Modeling Emissions by Scaling (TIMES) diurnal and day-of-week scaling (Nassar et al., 2013). The  $1^\circ \times 1^\circ$  uncertainty map is based on the combination of the global level FF uncertainty (one-sigma of 4.2%, Andres et al., 2014) and the grid level emission differences due to the different disaggregation methods (Oda et al., 2015). Note that these FF uncertainties are not considered in the inversions used for this product development.

Country-level fossil fuel emission estimates are obtained by aggregating the  $1^\circ \times 1^\circ$  estimates using the country mask.  
245 Uncertainties on country-level estimates are calculated using the fractional uncertainties of Andres et al. (2014).

#### 3.2 Net Carbon Exchange (NCE) and Net Biosphere Exchange (NBE)

We employ results from the v10 OCO-2 MIP, which is an international collaboration of atmospheric  $\text{CO}_2$  inversion modelers that produces ensembles of  $\text{CO}_2$  surface–atmosphere flux estimates by assimilating space-based OCO-2 retrievals of  $X_{\text{CO}_2}$  and in situ  $\text{CO}_2$  measurements. The v10 OCO-2 MIP is updated from the v9 OCO-2 MIP described in Peiro et al. (2022). Updates  
250 to the v10 OCO-2 MIP are presented here with additional details available at [https://gml.noaa.gov/ccgg/OCO2\\_v10mip/](https://gml.noaa.gov/ccgg/OCO2_v10mip/).



**Table 1.** Inversions specifications for each v10 OCO-2 MIP ensemble member.

Simulation name (reference)	Transport model	Driving meteorology	Meteorology resolution (degrees)	Prior NEE	Prior air-sea	Prior fire	Inverse Method
AMES (Philip et al., 2019, 2022)	GEOS-Chem	MERRA-2	4° × 5°	CASA-GFED4.1s	CT2019OI	GFEDv4.1s	4D-Var
Baker (Baker et al., 2006b, 2010)	PCTM	MERRA-2	1° × 1.25° prior, 4° × 5° opt	CASA-GFED3	Landschützer v4.4	GFEDv4	4D-Var
CAMS (Chevallier et al., 2005b)	LMDz	ERA5	1.9° × 3.75°	ORCHIDEE (climatological)	CMEMS	GFED4.1s	Variational
CMS-Flux <sup>c</sup> (Liu et al., 2021a)	GEOS-Chem	MERRA-2	4° × 5°	CARDAMOM	MOM-6	CARDAMOM	4D-Var
COLA (Liu et al., 2021b)	GEOS-Chem	MERRA-2	4° × 5°	VEGAS	Rödenbeck 2021	VEGAS	EnKF
CSU	GEOS-Chem	MERRA-2	4° × 5°	SiB-4 w/ MERRA-2	Landschützer v18	GFED4.1s	synthesis
CT (Jacobson et al., 2020)	TM5	ERA5	2° × 3°/ 1° × 1°	CT2019 CASA GFED4.1s	CT2019OI	CT2019 CASA GFED4.1s	EnKF
JHU <sup>a</sup> (Miller et al., 2020) (Chen et al., 2021b, a)	GEOS-Chem	MERRA-2	4° × 5°	CASA-GFED 4.1s	Takahashi	GFED4.1s	geostatistical/ 4D-Var
LoF <sup>b</sup> (Weir et al., 2021)	GEOS GCM	GEOS CGM	0.5° × 0.625°	CASA-GFED3	LoFI Takahasi	QFED	N/A
NIES (Maksyutov et al., 2021)	NIES-TM/ FLEXPART	ERA-5/ JRA-55	3.75° × 3.75°/ 0.1° × 0.1°	Zeng 2020	Landschützer 2020	GFAS	4D-Var
OU	TM5	ERA-Interim	4° × 6°	CASA-GFED3	Takahashi	GFEDv3	4D-Var
TM5-4DVar	TM5	ERA-Interim	2° × 3°	SiB-CASA	CT2019 Opt Clim	GFEDv4	4D-Var
UT (Deng et al., 2014, 2016)	GEOS-Chem	GEOS-FP	4° × 5°	BEPS	Takahashi	GFEDv4	4D-Var
WOMBAT (Zammit-Mangion et al., 2022)	GEOS-Chem	MERRA-2	2° × 2.5°	SiB-4 w/ MERRA-2	Landschützer 2020	GFED4.1s	Synthesis with MCMC

<sup>a</sup>Due to time constraints, this ensemble member is not included in this product; <sup>b</sup>This is not a traditional inversion and is not included in this product.



The v10 OCO-2 MIP consists of a number of inversion systems that perform a set of experiments following a standard protocol. Here, we include fluxes from 11 of the 14 MIP models (Table 1; CMS-Flux and JHU were excluded due to time constraints and LoFI was excluded because it employs a non-traditional inversion approach that does not follow the MIP protocol). There are five v10 OCO-2 MIP experiments that each ensemble member performs, which differ by the data that is  
255 assimilated (CO<sub>2</sub> datasets described in Sect. 3.2.1):

- **IS**: assimilates in situ CO<sub>2</sub> mole fraction measurements from an international observational network,
- **LNLG**: ACOS v10 land nadir and land glint total column dry-air mole fractions ( $X_{\text{CO}_2}$ ) from OCO-2,
- **LNLGIS**: assimilates both in situ and ACOS v10 OCO-2 land nadir and glint  $X_{\text{CO}_2}$  retrievals together,
- **OG**: assimilates ACOS v10 OCO-2 ocean glint  $X_{\text{CO}_2}$  retrievals
- 260 – **LNLGOGIS**: assimilates all the above datasets together.


For each experiment, each inversion group imposes a common fossil fuel emission dataset identical to the one described in Sect. 3.1. All other prior flux estimates were chosen independently by each modeling group and are listed in Table 1. The inversions assimilate the standardized v10 OCO-2 and in situ data from 6 September 2014 through 31 March 2021 (see  
265 Sect 3.2.1), with the length of spin-up period and in situ data assimilated during that period being left up to the discretion of each group in the MIP. Each modeling group submitted net air–sea fluxes and NBE across 2015–2020, interpolated from the native resolution to a 1° × 1° spatial grid at monthly resolution, which are publicly available for download from [https://gml.noaa.gov/ccgg/OCO2\\_v10mip/](https://gml.noaa.gov/ccgg/OCO2_v10mip/).

The performance of each atmospheric CO<sub>2</sub> inversion was evaluated through comparisons of the posterior CO<sub>2</sub> mole-fraction field (i.e., CO<sub>2</sub> fields simulated forward with the posterior fluxes) against independent in situ CO<sub>2</sub> measurements and OCO-2  
270  $X_{\text{CO}_2}$  retrievals that were withheld from the assimilation for validation, as well as  $X_{\text{CO}_2}$  retrievals from the Total Column Carbon Observing Network (TCCON; Wunch et al., 2011). The evaluation of the experiments is presented in Sect. 4, with additional analysis available from the v10 OCO-2 MIP website.

For this study, the best estimate of NCE is taken to be the ensemble median for each experiment (denoted  $\overline{\text{NCE}}_{\text{experiment}}$ )  
(. The uncertainty in NCE is calculated as an estimate of the standard deviation (denoted  $\sigma_{\text{NCE}}$ ) using the interquartile range  
275 (IQR) of the flux-inversion ensemble:

$$\sigma_{\text{NCE}} = \frac{\text{IQR}(\text{NCE})}{1.35}. \quad (4)$$

For country-level fluxes, the NCE estimates are first aggregated to country totals for each ensemble member before calculating the median and standard deviation. This is done because there are spatial covariances between 1° × 1° grid cells. Thus, first  
280 aggregating regions for each ensemble member accurately propagates the aggregate differences between regions across the ensemble members.



**Figure 3.** Assimilated observations for IS and LNLG v10 MIP experiments. Number of monthly (a) in situ CO<sub>2</sub> measurements and (b) ACOS v10 OCO-2 land nadir and land glint X<sub>CO<sub>2</sub></sub> retrievals binned into 10 s averages, and (c) ACOS v10 OCO-2 ocean glint X<sub>CO<sub>2</sub></sub> retrievals binned into 10 s averages. Spatial distribution of (d) in situ (e) ACOS v10 OCO-2 land X<sub>CO<sub>2</sub></sub> retrievals, and (f) ACOS v10 OCO-2 ocean X<sub>CO<sub>2</sub></sub> retrievals over 2015–2020. Shipboard and aircraft in situ CO<sub>2</sub> measurements are aggregated to a 2° × 2° spatial grid, surface site measurements are shown as scattered points, and ACOS v10 OCO-2 X<sub>CO<sub>2</sub></sub> retrievals are shown aggregated to a 2° × 2° spatial grid.

The NBE estimate is calculated by subtracting the ODIAC Fossil Fuel emissions from NCE. The variance in NBE is then taken to be the sum of the variances of NCE and FF:

$$\sigma_{\text{NBE}}^2 = \sigma_{\text{NCE}}^2 + \sigma_{\text{FF}}^2 \quad (5)$$

### 3.2.1 Atmospheric CO<sub>2</sub> data included in v10 OCO-2 MIP

285 In situ CO<sub>2</sub> measurements (Fig. 3a,d) are drawn from five data collections made available in Obspack format (Masarie et al., 2014). Those source ObsPacks and their references are listed in Table 2. The majority of data are from the openly available GLOBALVIEW+ program, but with some additional provisional data for 2020–21, and data from other programs not participating in the GLOBALVIEW+ project. CO<sub>2</sub> measurements are broadly divided into two categories: those measurements we identify as suitable for assimilation, and other measurements not suitable for assimilation.

290 In CO<sub>2</sub> inverse analyses, uncertainties ascribed to in situ measurements are a combination of the uncertainty in the measurement and a representativeness error from the forward model inability to accurately simulate the measurement (due to aspects like a coarse model grid). To characterize the representativeness error, we used an empirical scheme based on simulations from the v7 OCO-2 MIP (Crowell et al., 2019). In situ CO<sub>2</sub> measurements are simulated in a forward simulation, and then the model-data mismatch statistics are calculated to characterize the representativeness errors at each measurement location and  
 295 for each season. Although this was the standard method for characterizing uncertainties for modeled in situ measurements, each v10 OCO-2 MIP group was free to choose how to set the uncertainties in their specific set-ups.

Of the in situ measurements designated as being appropriate for assimilation, about 5% were withheld for cross-validation purposes. These data were chosen to be as independent as possible from the measurements that were assimilated. For quasi-continuous measurements, such as those taken every 15 minutes at NOAA tall towers, measurements were withheld for entire



**Table 2.** In situ CO<sub>2</sub> measurement collections used in the v10 OCO-2 MIP, with the total number of measurements between 6 September 2014 and 1 January 2021, and the numbers of measurements assimilated and withheld for cross-validation in the same period. More than 95% of the in situ measurements come from the GLOBALVIEW+ and CO<sub>2</sub> NRT ObsPacks, both of which are publicly available at <https://gml.noaa.gov/ccgg/obspack/data.php>.

ObsPack name	Total No. Measurements	Assimilated	Withheld	Reference
obspack_CO <sub>2</sub> _1_GLOBALVIEWplus_v6.1_2021-03-01	9,611,095	766,179	38,483	Schuldt et al. (2021b)
obspack_CO <sub>2</sub> _1_NRT_v6.1.1_2021-05-17	755,477	62,011	2,996	Schuldt et al. (2021a)
obspack_CO <sub>2</sub> _1_NIES_Shipboard_v3.0_2020-11-10	418,496	216,963	12,766	Tohjima et al. (2005); Nara et al. (2017)
obspack_CO <sub>2</sub> _1_AirCore_v4.0_2020-12-28	55,620			Baier et al. (2021)
obspack_multi-species_1_manauis_profiles_v1.0_2021-05-20	3,194			Miller et al. (2021)
<b>TOTAL</b>	<b>10,843,882</b>	<b>1,045,153</b>	<b>54,245</b>	



300 days: we chose 5% of the days in the dataset, and we withheld every assimilable measurement on that day. This is also how CO<sub>2</sub> measurements on NIES ships were treated. Entire aircraft profiles in the NOAA light-aircraft profiling network are assumed to consist of vertically correlated measurements, so entire profiles were withheld: we chose 5% of aircraft profiles to withhold. Most flask sites have measurement sampling protocols intended to ensure independence; they are often taken at weekly or biweekly intervals during meteorological conditions meant to allow regional background air masses to be sampled. Thus, we  
305 chose to withhold 5% of assimilable flask measurements. We also verified that datasets at the same site were withheld on the same days; aircraft profiles over tower sites were, for instance, withheld on the same days that tower data were withheld.

OCO-2 land (Fig. 3b,e) and ocean (Fig. 3c,f) X<sub>CO<sub>2</sub></sub> retrievals are performed using version 10 of NASA's Atmospheric CO<sub>2</sub> Observations from Space (ACOS) full-physics retrieval algorithm (O'Dell et al., 2018). A common set of OCO-2 retrieval "super-obs" data were derived from these retrievals and were assimilated by each modeling group. These "super-obs" are  
310 obtained by aggregating retrievals into 10 s averages (which better match the coarse transport models grid cells used in the inversions) following the same procedure as the v9 OCO-2 MIP (Peiro et al., 2022). Specifically, individual scenes within the 10 s span are weighted according to the inverse of the square of the X<sub>CO<sub>2</sub></sub> uncertainty (standard deviations) produced by the retrieval, and correlations of +0.3 for land scenes and +0.6 for ocean scenes are assumed when calculating the uncertainty on the 10-second averages (see Sect. 3.2.1 of Baker et al., 2022); transport model errors are also considered (based on Schuh  
315 et al., 2019). Only 10 s spans with 10 or more good quality retrievals were used (sparser data being thought to be more prone to cloud-related biases). In the same vein as was done for the in situ data, X<sub>CO<sub>2</sub></sub> data from 5% of the orbits (entire orbits were withheld), chosen at random, were withheld for evaluation purposes.

### 3.3 Country-level $F_{\text{rivers export}}$

Rivers transport carbon laterally across land regions (e.g., to a lake) and from the land to the ocean. This lateral transport must  
320 be accounted for to quantify the total change in terrestrial carbon in a given region. However, there is considerable uncertainty in lateral carbon flux by rivers. To account for this, we use two independent estimates of country-level totals: one from the Dynamic Land Ecosystem Model (DLEM), and the other based on Deng et al. (2022) who use the Global NEWS model (Mayorga et al., 2010) and observations across COastal Segmentation and related CATchments (COSCATs) (Meybeck et al., 2006) that include DIC (of atmospheric origin), DOC and POC. These datasets cover 2015–2019. For 2020, we impose the  
325 2015–2019 mean.

The DLEM is a process-based terrestrial ecosystem model that couples biophysical, soil biogeochemical, plant physiological and riverine processes with vegetation and land-use dynamics to simulate and predict the vertical fluxes, lateral fluxes, and storage of water, carbon, GHGs, and nutrient dynamics in terrestrial ecosystems and their interfaces with the atmosphere and land-ocean continuum (Tian et al., 2010, 2015a). There are three major processes involved in simulating the export of water,  
330 carbon, and nutrients from land surface to the coastal ocean: 1) the generation of runoff and leachates, 2) the leaching of water, carbon and nutrients from land to river networks in the form of overland flow and base flow, and 3) transport of riverine materials along river channels from upstream areas to coastal regions. The key processes and parameterization in the DLEM have been described in previous publications regarding the water discharge (Liu et al., 2013; Tao et al., 2014), riverine carbon





fluxes (Ren et al., 2015, 2016; Tian et al., 2015b; Yao et al., 2021), and riverine nitrogen fluxes (Yang et al., 2015; Tian et al.,  
335 2020) from the terrestrial ecosystem to coastal oceans. The newly improved DLEM aquatic module better addresses processes  
within global small streams, which were recognized as hotspots of GHG emissions (Yao et al., 2020, 2021). DLEM produces  
estimates of the land loadings of carbon species (DIC, DOC, and POC), CO<sub>2</sub> degassing and carbon burial during transporting,  
and the exports of carbon (DIC, DOC, and POC) to the ocean for 105 basin-level segmentations (modified from COSCATs)  
(Meybeck et al., 2006). To estimate country totals, we map the basin carbon loss across land by assuming that the net carbon  
340 flux occurs uniformly across each basin. We then use the country mask to estimate the country totals for each region.

Deng et al. (2022) estimate the lateral carbon export by rivers to the coast minus the imports from rivers entering in each  
country (for relevant cases), including DOC, POC and DIC of atmospheric origin. Estimates of DOC, POC and DIC are  
obtained from the Global NEWS model (Mayorga et al., 2010), with a correction based on Resplandy et al. (2018) so that the  
global total exported to the coastal ocean is 2.86 PgCO<sub>2</sub> yr<sup>-1</sup> (0.78 PgC yr<sup>-1</sup>). Deng et al. (2022) perform a correction to the  
345 Global NEWS estimates to remove the contribution of lithogenic carbon, using the methodology of Ciais et al. (2021).

For the analysis that follows, we estimate country-level totals of riverine lateral carbon fluxes by combining the estimates of  
DLEM with those of Deng et al. (2022). We take the mean of the two estimates to be the best estimate and take the magnitude  
of the difference between the estimates to be the one-sigma uncertainty. Figure S1 shows the 2015–2019 mean annual net  
riverine lateral carbon fluxes. Fluxes are uniformly negative, implying a net flux of carbon from the land to the ocean and  
350 reduction in stored carbon for all countries. Fluxes are most negative in tropical rainforest and tropical monsoon climates, and  
they are smallest in more arid regions.

### 3.4 Country-level $F_{\text{wood trade}}$ and $F_{\text{crop trade}}$

Wood and crop products are traded between nations. We estimate the annual lateral fluxes of carbon due to this trade following  
the approaches of Deng et al. (2022) and Ciais et al. (2021). This approach utilizes crop and wood trade data compiled by  
355 the Food and Agriculture Organization of the United Nations (FAO, <http://www.fao.org/faostat/en/#data>). The crop flux was  
estimated from the annual trade balance of 171 crop commodities calculated for each country. For wood products, we use the  
bookkeeping model of Mason Earles et al. (2012) to calculate the fraction of imported carbon in wood products that is oxidized  
in each of 270 countries during subsequent years. One-sigma uncertainties in country-level fluxes are assumed to be 30% of  
the mean value. This dataset covers 2015–2019. For 2020, we assume fluxes equal to the 2015–2019 mean. The net crop and  
360 wood lateral fluxes and their uncertainties are shown in Fig. S2.

### 3.5 1° × 1° lateral flux estimates

Lateral fluxes at a higher resolution (1° × 1°) follow similar principles to national values but were estimated separately with  
different implementation choices. High-resolution proxy data (satellite-derived NPP, population or livestock maps, etc.) enabled  
subnational disaggregation. For each 1° × 1° grid cell, we assume the standard deviation of the mean flux to be 30% for  
365  $F_{\text{wood trade}}$  and  $F_{\text{crop trade}}$ , and 60% for  $F_{\text{rivers export}}$ .



### 3.6 Estimate of carbon stock loss ( $\Delta C_{\text{loss}}$ )

Finally, we calculate  $\Delta C_{\text{loss}}$  using Eqn. 2 with the datasets described above. Assuming that the components contributing to  $\Delta C_{\text{loss}}$  are independent, we calculate the uncertainty on  $\Delta C_{\text{loss}}$  by combining the uncertainties (one standard deviations) from the component fluxes in quadrature:

$$370 \quad \sigma_{\Delta C_{\text{loss}}}^2 = \sigma_{\text{NBE}}^2 + \sigma_{F_{\text{crop trade}}}^2 + \sigma_{F_{\text{wood trade}}}^2 + \sigma_{F_{\text{rivers export}}}^2 \quad (6)$$

## 4 Evaluation of v10 OCO-2 MIP experiments


The performance of top-down  $\text{CO}_2$  flux estimates can be impacted by a number of factors, including biases in the assimilated data, model transport, prior constraints, and in the inversion architectures. Therefore, evaluating the performance of v10 OCO-2 MIP fluxes against independent datasets is critical for assuring high quality flux estimates. Here, we evaluate the v10 OCO-2 MIP experiments in two ways. First, we compare the posterior  $\text{CO}_2$  fields against independent  $\text{CO}_2$  measurements (Sect. 4.1).  
375 Second, we compare the inferred air–sea  $\text{CO}_2$  flux against estimates based on surface ocean  $\text{CO}_2$  partial pressure ( $p\text{CO}_2$ ) measurements (Sect. 4.2).

### 4.1 Evaluation of posterior $\text{CO}_2$ fields

We consider four atmospheric  $\text{CO}_2$  datasets:

- 380 1. Withheld in situ  $\text{CO}_2$  measurements. These are measurements contained in the Obspack collection described in Sect. 3.2.1 but intentionally withheld for evaluation purposes. Independence from the assimilated data is ensured following the steps described in Sect. 3.2.1.
- 385 2.  $X_{\text{CO}_2}$  retrievals from the TCCON. These data are acquired from a network of ground-based Fourier Transform Spectrometers measuring direct solar spectra from which  $X_{\text{CO}_2}$  is retrieved (Wunch et al., 2011). For this analysis, we include 30 TCCON sites listed in table A1. These data are filtered and aggregated following the method outlined in Appendix C of Crowell et al. (2019).
3. Withheld OCO-2 land glint and land nadir  $X_{\text{CO}_2}$  retrievals. These data could have been assimilated, but they are intentionally withheld for evaluation purposes (Sect. 3.2.1).
- 390 4. Withheld OCO-2 ocean glint  $X_{\text{CO}_2}$  retrievals. These data could have been assimilated, but they are intentionally withheld for evaluation purposes (Sect. 3.2.1).


We first perform a simple check on the inversion results by comparing the atmospheric  $\text{CO}_2$  growth rate estimated from the v10 OCO-2 MIP experiments to that derived directly from NOAA  $\text{CO}_2$  measurements (Fig. 4). The growth rate is estimated from  $\text{CO}_2$  measurements and model co-samples at “marine boundary layer” sites, which predominantly observe well-mixed marine boundary layer air representative of a large volume of the atmosphere. A smooth curve is then fit to these data to estimate



**Figure 4.** Mean 2015–2019 global mean CO<sub>2</sub> growth rate estimated from NOAA site measurements and for the v10 OCO-2 MIP experiments. The estimates of the CO<sub>2</sub> growth rate for each experiment are computed by sampling the model CO<sub>2</sub> fields at the same times and locations as those used to derive the NOAA measurement-based estimate. Each v10 OCO-2 MIP experiment is shown as a box plot, with the error bars showing the full range, the shaded region showing the interquartile range, and the solid line showing the median ensemble member of the ensemble.


395 the global growth rate (Thoning et al., 1989). This is the same method employed by NOAA to report the CO<sub>2</sub> growth rate  
(gml.noaa.gov/ccgg/trends/). We estimate the uncertainty in the measurement-based growth rate from the difference between  
the growth rate estimated here and that reported on the NOAA website. Differences between these estimates are primarily  
driven by differences in measurement sampling used for the website relative to that used here (as we are limited to withheld  
co-samples here). We calculate the uncertainty as the standard error of the mean for the differences between the growth rates  
400 estimated here and by NOAA across 2015–2019. This gives an uncertainty on the 5-year growth rate of  $\pm 0.053$  ppm yr<sup>-1</sup>.  
Note that NOAA reports the growth rate using the X2019 scale, whereas our estimates here are from the X2007 scale, which  
may contribute to the differences. We find that the IS, LNLG, and LNLGIS experiments show good agreement with the NOAA  
estimate over this period. However, both the OG and LNLGOGIS experiments are found to have a high bias. This suggests that  
there may be a spurious trend in the v10 OCO-2 ocean glint X<sub>CO<sub>2</sub></sub> retrievals of 0.04–0.13 ppm yr<sup>-1</sup> (OG experiment bias) that  
405 impacts flux estimates from both experiments that assimilate ocean glint data.

Second, we estimate the overall data–model agreement as the root-mean-square (RMS) error for the the withheld in situ  
CO<sub>2</sub>, TCCON X<sub>CO<sub>2</sub></sub>, withheld OCO-2 land X<sub>CO<sub>2</sub></sub>, and withheld OCO-2 ocean X<sub>CO<sub>2</sub></sub> (Fig. 5). For the in situ and OCO-2  
data, the normalized RMS is shown, meaning that the data–model difference is divided by the observational uncertainty (one-  
sigma). Overall, we find reasonably good agreement between the evaluation datasets and posterior fields for all experiments.  
410 The OG experiment gives the largest RMS errors against the withheld in situ CO<sub>2</sub>, TCCON X<sub>CO<sub>2</sub></sub>, and OCO-2 land X<sub>CO<sub>2</sub></sub>.  
This provides further evidence that the ocean glint data may have some residual biases that adversely impact the flux estimates.



**Figure 5.** 2015–2020 root-mean-square (RMS) error between the v10 OCO-2 MIP experiments and (a) TCCON  $X_{CO_2}$  retrievals, (b) withheld in situ  $CO_2$  measurements, (c) withheld OCO-2 land  $X_{CO_2}$  retrievals, and (d) withheld OCO-2 ocean  $X_{CO_2}$  retrievals. For the comparisons with withheld in situ and OCO-2 observations, the normalized RMS estimate is plotted (that is, the data–model mismatch is divided by the observational uncertainty). Note that NIES IS and CSU co-samples are not available and not included in this plot.

Finally, we examine the mean bias over 2015–2020 for  $30^\circ$  latitude bins (Fig. 6). Similar to previous comparisons, we find that the OG experiment stands out as being more biased against the independent data relative to the other experiments. In particular, the data–model difference for the OG experiment tends to be low (higher modeled  $CO_2$ ) than the evaluation datasets. This is particularly evident in the northern extratropics. Over  $30^\circ$ – $60^\circ$  N, where independent data is densest, we find that the OG ensemble median is biased by  $-0.69$  ppm against TCCON,  $-0.74$  ppm against withheld in situ, and  $-0.48$  ppm against withheld OCO-2 LNLG, suggesting a possible meridional bias in the OCO-2 ocean  $X_{CO_2}$  retrievals. The IS, LNLG, and LNLGIS experiments tend to show similar data–model differences, suggesting limited ability to distinguish between the performance of these inversions in large-scale features.



**Figure 6.** Median bias (data minus model) over 30° latitude bins averaged over 2015–2020 for (a) TCCON  $X_{CO_2}$  retrievals, (b) withheld in situ  $CO_2$  measurements, (c) withheld OCO-2 land  $X_{CO_2}$  retrievals, and (d) withheld OCO-2 ocean  $X_{CO_2}$  retrievals. Note that and NIES IS and CSU co-samples are not available and not included in this plot.

All experiments show some biases against TCCON sites. In particular, low biases (high modeled  $CO_2$ ) are found for 0°–30° S and 60°–90° N. The underlying cause for these differences is unknown. Figure S3 shows the monthly-mean data–model differences for each TCCON site and each experiment. The differences can be quite variable between sites, but are generally similar between experiments (for IS, LNLG, and LNLGIS). Some of these differences may be related due to representativeness errors, particularly for urban sites. For example, Caltech and JPL are within Los Angeles County and show a large positive bias, while nearby Edwards is less impacted by urban emissions and shows a much smaller bias (Schuh et al., 2021). However, other differences are harder to explain, such as a negative trend in the data–model bias for Park Falls and positive at Darwin during the 2015–2020 period. Site-to-site biases among TCCON sites may also contribute to these differences.

Overall, this analysis finds that the OG experiment shows the poorest agreement against the evaluation datasets (excluding the withheld ocean glint data). The LNLGOGIS experiment shows the second worst performance against evaluation datasets, while the remaining experiments (IS, LNLG, and LNLGIS) all show good agreement against the evaluation data. These results suggest that there may be residual biases in the OCO-2 ocean glint dataset that adversely impact the OG and LNLGOGIS experiments.



## 4.2 Comparison of air–sea fluxes with pCO<sub>2</sub>-based estimates

435 The exchange of CO<sub>2</sub> between the atmosphere and the ocean (air–sea flux) can be estimated from measurements of the surface  
ocean partial pressure of CO<sub>2</sub> (pCO<sub>2</sub>). These pCO<sub>2</sub> data are extrapolated to global maps and combined with gas transfer  
velocity parameterizations to infer global maps of the air–sea CO<sub>2</sub> fluxes (Fay et al., 2021). Although significant uncertainties  
remain, particularly in accurately representing the gas transfer velocity (Fay et al., 2021), comparisons between the pCO<sub>2</sub>-  
based air–sea fluxes and v10 OCO-2 MIP experiments can inform possible biases between estimates and inform potential areas  
440 for future research.


Here, we compare v10 OCO-2 MIP air–sea fluxes to an ensemble of air–sea flux estimates from SeaFlux (Fay et al., 2021;  
Gregor and Fay, 2021). SeaFlux developed a standardized approach to harmonize and extend six air–sea CO<sub>2</sub> flux products  
from as many surface pCO<sub>2</sub> products: JENA-MLS (Rödenbeck et al., 2013), MPI-SOMFFN (Landschützer et al., 2014, 2020),  
CMEMS-FFN (Denvil-Sommer et al., 2019; Chau et al., 2022), CSIR-ML6 (Gregor et al., 2019), JMA-MLR (Iida et al.,  
445 2021), and NIES-FNN (Zeng et al., 2014). For each pCO<sub>2</sub> product, we examine the mean of three air–sea fluxes obtained  
using different wind reanalysis datasets to estimate the gas transfer parameterization (ERA5, JRA55, and CCMP2). The spread  
among these six estimates provides a measure of uncertainty in the extrapolation of pCO<sub>2</sub> data to a global grid, but does not  
account for errors in the gas transfer velocity formulation nor the uncertainties in the reanalysis winds used as input (Fay et al.,  
2021). Note that the prior estimates of air–sea CO<sub>2</sub> fluxes in v10 OCO-2 MIP experiments are generally pCO<sub>2</sub>-based flux  
450 estimates, and therefore not independent from the SeaFlux datasets.

Figure 7 shows the 2015–2019 mean air–sea fluxes for each of the six SeaFlux products and for the v10 OCO-2 MIP exper-  
iments across 30° latitude bands and large ocean regions. Over the global ocean, the pCO<sub>2</sub>-based air–sea fluxes tend to give  
stronger removals (median =  $-10.0 \text{ PgCO}_2 \text{ yr}^{-1}$  or  $-2.7 \text{ PgC yr}^{-1}$ , range =  $-0.2$  to  $-12.9 \text{ PgCO}_2 \text{ yr}^{-1}$  or  $-3.5$  to  $-2.5 \text{ PgC yr}^{-1}$ )  
than the v10 OCO-2 MIP, which range from  $-7.9 \pm 1.9 \text{ PgCO}_2 \text{ yr}^{-1}$  ( $-2.1 \pm 0.5 \text{ PgC yr}^{-1}$ ) for the IS experiment to  
455  $-10.2 \pm 1.28 \text{ PgCO}_2 \text{ yr}^{-1}$  ( $-2.8 \pm 0.4 \text{ PgC yr}^{-1}$ ) for the OG experiment. On regional scales, the v10 OCO-2 MIP exper-  
iments overlap with the pCO<sub>2</sub>-based estimates except for the northern high latitudes (60°–90° N), where pCO<sub>2</sub>-based estimates  
suggest a systematically larger removals. Similarly, the pCO<sub>2</sub>-based estimates tend to give greater removals over the southern  
midlatitudes (20°–50° S).

The different v10 OCO-2 MIP experiments tend to give similar air–sea fluxes, except for the OG experiment in the tropics.  
460 Although not systematic, the OG experiment suggests weaker emissions in the tropics of  $0.2 \pm 1.3 \text{ PgCO}_2 \text{ yr}^{-1}$  ( $0.05 \pm 0.34 \text{ PgC yr}^{-1}$ )  
relative to the median pCO<sub>2</sub>-based estimate of  $1.6 \text{ PgCO}_2 \text{ yr}^{-1}$  ( $0.43 \text{ PgC yr}^{-1}$ ) with a range of  $0.4$  to  $1.8 \text{ PgCO}_2 \text{ yr}^{-1}$   
( $0.10$  to  $0.50 \text{ PgC yr}^{-1}$ ). Thus, similar to the evaluation of posterior CO<sub>2</sub> fields, the OG experiment is an outlier among the v10  
OCO-2 MIP experiments, further supporting the possibility that residual biases may exist in the ocean glint X<sub>CO<sub>2</sub></sub> retrievals.

## 5 Metrics for interpreting country flux estimates

465 To aid users in interpreting top-down country-level flux estimates, we provide two metrics. The first metric is called the “Z  
statistic” and quantifies the statistical agreement between the IS and LNLG NCE estimates, and thus gives an indication of



**Figure 7.** (a) Zonal-mean air–sea CO<sub>2</sub> flux (positive values represent flux towards atmosphere) for 30° increments of latitude based on 1° × 1° estimates averaged over 2015–2019. (b) air–sea CO<sub>2</sub> flux for six large ocean regions. Colored bars show the MIP experiment results (median +/- one standard deviation) and the symbols show the pCO<sub>2</sub>-based air–sea fluxes from the six SeaFlux products.

how robust flux estimates are across the v10 OCO-2 MIP experiments (Sect. 5.1). The second metric is called the Fractional Uncertainty Reduction (FUR) and informs the impact of the assimilated CO<sub>2</sub> data on the estimated fluxes (Sect. 5.2).


### 5.1 Z statistic

470 The Z statistic is defined as,

$$\text{Z statistic} = \frac{\overline{\text{NCE}}_{\text{LNLG}} - \overline{\text{NCE}}_{\text{IS}}}{\text{std}(\text{NCE}_{\text{LNLG}} - \text{NCE}_{\text{IS}})}, \quad (7)$$

where the denominator represents the standard deviation in  $\text{NCE}_{\text{LNLG}} - \text{NCE}_{\text{IS}}$  across the ensemble members. Differences in NCE and  $\Delta C_{\text{loss}}$  between v10 OCO-2 MIP experiments can be considerable. As an example, Fig. 8a shows that differences between  $\overline{\text{NCE}}_{\text{LNLG}}$  and  $\overline{\text{NCE}}_{\text{IS}}$  are notable for South America and Africa. The LNLG experiment gives more positive  $\Delta C_{\text{loss}}$  (carbon loss from land) over northern sub-Saharan Africa and northeast South America, but more negative  $\Delta C_{\text{loss}}$  over southern tropical Africa, southern and eastern South America, and southeast Asia. We examine the Z statistic (Fig. 8b) to quantify the statistical significance of these difference (magnitude greater than 1.96 indicates statistically significant differences at level  $\alpha = 0.05$ ). Most countries do not have statistically significant differences, indicating relatively good agreement between the IS

475



**Figure 8.** Difference between LNLG and IS experiments. (a)  $\overline{\text{NCE}}_{\text{LNLG}}$  minus  $\overline{\text{NCE}}_{\text{IS}}$ , and (b) The Z statistic (Eqn. 7) indicating the difference between LNLG and IS experiments.

and LNLG ensembles. Significant differences primarily occur in small to mid-sized tropical countries. Canada also shows a  
480 systematic difference driven by small uncertainties in the IS and LNLG estimates.


## 5.2 Fractional uncertainty reduction (FUR)

Byrne et al. (2022) reports the uncertainty in NCE as the standard deviation across v10 OCO-2 MIP ensemble members  
(estimated using Eqn. 4). This metric incorporates uncertainties related to model transport and aspects of the inversion config-  
uration, such as optimization technique and a priori flux estimates. However, this metric is different to the uncertainty metric  
485 usually computed in a Bayesian framework, that is, the Bayesian posterior uncertainty. That uncertainty quantifies the impact  
of errors in the observations and prior constraints on the posterior flux estimates. The Bayesian posterior uncertainty is not  
reported for practical reasons, as the majority of contributing models do not calculate this quantity, so it is not possible to  
calculate this quantity across the ensemble.

In this section, we examine the posterior uncertainty estimates from two contributing inversion systems (CAM5 and TM5-  
490 4DVar) and compare these estimates to the ensemble-based uncertainty estimate provided with the dataset. Then, we define  
the metric of Fractional Uncertainty Reduction (FUR) between the posterior and prior NCE estimates based on the TM5-  
4DVar model (as CAM5 does not estimate uncertainties for the LNLGIS and LNLGOGIS experiments), which can be used to  
understand the relative impact of assimilated atmospheric  $\text{CO}_2$  data on estimates of country-level NCE and  $\Delta C_{\text{loss}}$ .

Both CAM5 and TM5-4DVar estimate  $\text{CO}_2$  fluxes using four-dimensional variational assimilation (4D-Var) and estimate  
495 posterior uncertainty estimates using a Monte Carlo method derived by Chevallier et al. (2007). The realism of the prior and  
posterior CAM5 uncertainty estimates have already been the topic of several studies (see Chevallier, 2021, and references  
therein). Figure 9 shows the ensemble-based uncertainty, prior/posterior uncertainty from CAM5 (prior, IS and LNLG only)  
and prior/posterior uncertainty from TM5-4DVar for four countries. Notably, the magnitudes of the prior/posterior uncertainties  
from CAM5 and TM5-4DVar are quite different, with CAM5 uncertainties being 2–8 times larger. Differences in prior/poste-






**Figure 9.** (top)  $\overline{NCE}$  and (bottom)  $\sigma_{NCE}$  for four countries in 2018. The v10 OCO-2 MIP ensemble spread-based error estimate is shown in black, the TM5-4DVar Bayesian uncertainty estimate is shown in red, and the CAMS Bayesian uncertainty estimate is shown in green (only for Prior, IS, and LNLG).

500 rior uncertainties of this magnitude are not unusual among inversion systems, and highlight the sensitivity of Bayesian uncertainty estimates to choices about prior uncertainties. Both CAMS and TM5-4DVar posterior uncertainties are smaller relative to their prior by similar amounts, driven by the assimilated CO<sub>2</sub> data. The magnitude of the ensemble-based uncertainty tends to fall in-between the CAMS and TM5-4DVar estimates. However, the CAMS and TM5-4DVar posterior uncertainty estimates decrease as more data are assimilated (as expected), while the ensemble spread does not. In fact, the ensemble spread increases  
 505 with data density in some cases (e.g., Australia LNLGIS). Thus, overall, we find that the ensemble-based uncertainty estimate is of similar magnitude to the prior/posterior estimate, but that the magnitude of posterior uncertainty is quite dependent on the assumed prior uncertainty.

We now calculate the FUR metric in NCE from the TM5-4DVar Bayesian uncertainties (note that we use TM5-4DVar only because CAMS does not report LNLGIS or LNLGOGIS uncertainties). FUR is calculated from the prior flux standard deviation  
 510 ( $\sigma_{\text{prior}}$ ) and posterior flux standard deviation ( $\sigma_{\text{posterior}}$ ) as:

$$\text{FUR} = 1 - \frac{\sigma_{\text{posterior}}}{\sigma_{\text{prior}}} \quad (8)$$

This quantity ranges between 0 and 1, with larger values indicating that the Bayesian uncertainties have decreased more (relative to the prior) due to the observational constraints from assimilated data. This metric is useful for understanding how the assimilation of data influences the NCE and  $\Delta C_{\text{loss}}$  estimates, which may not be captured by the ensemble spread. For  
 515 example, Saudi Arabia has a small NCE uncertainty estimate but this is largely driven by prior knowledge that biosphere CO<sub>2</sub> fluxes and the atmospheric CO<sub>2</sub> data has little impact on the NCE estimate.



**Figure 10.** Estimate of the Fractional Uncertainty Reduction (FUR) on the v10 OCO-2 MIP estimates for each experiment based on Bayesian uncertainty estimates from the TM5-4DVar inversion.

Figure 10 shows FUR for the IS, LNLG, LNLGIS, and LNLGOGIS experiments. FUR is larger in regions with denser observational coverage. For example, the IS FUR is close to 1 in the USA and parts of Europe, reflecting dense CO<sub>2</sub> measurements, but it remains small for many tropical countries, where sampling is sparse. Meanwhile, the LNLG experiment generally has larger FUR values than the IS experiment in the tropics, reflecting denser sampling, but has lower values for some small high-latitude countries, such as in Scandinavia.

## 6 Dataset description

The dataset described in this paper, Byrne et al. (2022), provides annual totals of country-level and  $1^\circ \times 1^\circ$  gridded  $\Delta C_{\text{loss}}$ , NBE, NCE,  $F_{\text{rivers export}}$ , and the combined  $F_{\text{crop trade}} + F_{\text{wood trade}}$  fluxes, as well as their uncertainties over 2015–2020. In addition, the country-level Z statistic (Eqn. 7) and FUR (Eqn. 8) metrics are provided to help interpret the flux and stockchange estimates. These data are provided for the v10 OCO-2 MIP IS, LNLG, LNLGIS, and LNLGOGIS experiments. The OG experiment is excluded due to poor evaluation against independent CO<sub>2</sub> measurements and pCO<sub>2</sub>-based air–sea fluxes, likely due to residual X<sub>CO<sub>2</sub></sub> biases in the OCO-2 ocean glint X<sub>CO<sub>2</sub></sub> retrievals (Sect. 4). We note that biases in ocean glint X<sub>CO<sub>2</sub></sub> retrievals will also adversely impact flux estimates from the LNLGOGIS, and caution against using these data when they show



530 differences from the IS, LNLG, and LNLGIS experiments. Future improvements to the OCO-2  $X_{\text{CO}_2}$  retrievals are expected to reduce residual  $X_{\text{CO}_2}$  biases and thus the quality of the LNLGOGIS experiment is expected to improve in future OCO-2 MIP experiments.

For the  $1^\circ \times 1^\circ$  gridded dataset, we emphasize that caution is needed in interpreting these data. As discussed in Sect. 1.3, atmospheric  $\text{CO}_2$  inversion analyses provide the best constraints on the largest spatial scales (e.g., continental-to-global). The confidence in these top-down estimates decreases at smaller spatial scales. The minimum spatial resolution for robust flux estimates is dependent on the density and precision of the measurements, and is challenging to quantify. However, scales smaller than France or Germany in geographic extent are unlikely to be meaningfully constrained. Thus, we recommend only using  $1^\circ \times 1^\circ$   $\text{CO}_2$  fluxes aggregated to larger spatial scales. In aggregating, we recommend propagating uncertainties by assuming first 100% correlation (sum of the  $1^\circ \times 1^\circ$  uncertainties) and then 0% correlation (square root of the sum of the squared uncertainties) between grid cells. We strongly encourage contacting the authors before using the gridded  $1^\circ \times 1^\circ$  dataset.

These data are available for download from the Committee on Earth Observation Satellites' (CEOS) website: <https://doi.org/10.48588/npf6-sw92>. The country-level data are available for download as comma-separated values (CSV), Network Common Data Form (NetCDF) and Microsoft Excel worksheet files. The  $1^\circ \times 1^\circ$  gridded dataset is available as a NetCDF file.

## 7 Characteristics of the dataset

Globally, over 2015–2020, we report FF emissions of  $35.79 \pm 1.50 \text{ PgCO}_2 \text{ yr}^{-1}$  ( $9.76 \pm 0.41 \text{ PgC yr}^{-1}$ ),  $F_{\text{rivers export}}$  of  $-3.35 \pm 0.59 \text{ PgCO}_2 \text{ yr}^{-1}$  ( $-0.91 \pm 0.16 \text{ PgC yr}^{-1}$ ), and globally balanced  $F_{\text{crop trade}}$  and  $F_{\text{wood trade}}$ . Table 3 gives the global annual mean changes in the atmospheric burden of  $\text{CO}_2$ ,  $\Delta C_{\text{gain}}$  and ocean sequestration. Across the experiments, the median fraction of fossil fuel emissions remaining in the atmosphere is 55–56%, while 32–36% is sequestered by the ocean and 9–13% is sequestered by terrestrial ecosystems. Note that this omits land use change (LUC) emissions of  $\sim 3.85 \text{ PgCO}_2 \text{ yr}^{-1}$  ( $\sim 1.05 \text{ PgC yr}^{-1}$ , Friedlingstein et al., 2022), which are compensated for by additional carbon uptake by land. Of the combined FF+LUC emissions, 50% remains in the atmosphere, 29–33% is sequestered by the ocean and 18–21% is sequestered by terrestrial ecosystems. Relative to Global Carbon Budget 2021 (GCB 2021; Friedlingstein et al., 2022) we find 2.24–3.53  $\text{PgCO}_2 \text{ yr}^{-1}$  ( $0.61\text{--}0.96 \text{ PgC yr}^{-1}$ ) less removal by land (mean/median difference) but greater removal by the ocean of 0.87–2.24  $\text{PgCO}_2 \text{ yr}^{-1}$  ( $0.24\text{--}0.61 \text{ PgC yr}^{-1}$ ), however, these difference are consistent within one standard deviation of the mean/median values. Interestingly, we report greater removals by the ocean than GCB 2021 but reduced air–sea flux relative to SeaFlux. This can be explained by the fact that  $\text{pCO}_2$ -based air–sea flux estimates generally give larger mean ocean carbon uptake than model estimates (Fay and McKinley, 2021) and that we estimate a larger  $F_{\text{rivers export}}$  than GCB 2021.

Meridionally, NCE is largest in the northern extratropics, coinciding with the largest FF emissions (Fig. 11). However, the northern extratropics also show negative  $\Delta C_{\text{loss}}$ , implying increasing terrestrial carbon stocks, particularly between  $30^\circ\text{--}60^\circ \text{ N}$ . NCE is less positive in the tropics, primarily due to lower FF emissions. However, this region tends to show neutral-to-positive




**Table 3.** 2015–2020 global mean atmospheric increase, terrestrial carbon gain ( $\Delta C_{\text{gain}}$ ) and ocean carbon gain from the IS, LNLG, LNLGIS, and LNLGOGIS experiments (mean/median  $\pm$  one standard deviation). Positive values of  $\Delta C_{\text{gain}}$  and ocean carbon gain indicate increases in carbon stocks. GCB 2021 were obtained from the Global Carbon Budget 2021 (Friedlingstein et al., 2022) with  $\Delta C_{\text{gain}}$  calculated as the difference between the land sink and land-use change emissions with errors propagated in quadrature.

Experiment	Atmosphere	$\Delta C_{\text{gain}}$	Ocean carbon gain
IS	$19.73 \pm 0.19 \text{ PgCO}_2 \text{ yr}^{-1}$ ( $5.38 \pm 0.05 \text{ PgC yr}^{-1}$ )	$4.58 \pm 2.44 \text{ PgCO}_2 \text{ yr}^{-1}$ ( $1.25 \pm 0.66 \text{ PgC yr}^{-1}$ )	$11.35 \pm 2.01 \text{ PgCO}_2 \text{ yr}^{-1}$ ( $3.10 \pm 0.55 \text{ PgC yr}^{-1}$ )
LNLG	$19.64 \pm 0.09 \text{ PgCO}_2 \text{ yr}^{-1}$ ( $5.36 \pm 0.02 \text{ PgC yr}^{-1}$ )	$3.29 \pm 2.93 \text{ PgCO}_2 \text{ yr}^{-1}$ ( $0.90 \pm 0.80 \text{ PgC yr}^{-1}$ )	$12.91 \pm 2.63 \text{ PgCO}_2 \text{ yr}^{-1}$ ( $3.52 \pm 0.72 \text{ PgC yr}^{-1}$ )
LNLGIS	$19.64 \pm 0.06 \text{ PgCO}_2 \text{ yr}^{-1}$ ( $5.36 \pm 0.02 \text{ PgC yr}^{-1}$ )	$4.19 \pm 2.77 \text{ PgCO}_2 \text{ yr}^{-1}$ ( $1.14 \pm 0.75 \text{ PgC yr}^{-1}$ )	$11.98 \pm 2.32 \text{ PgCO}_2 \text{ yr}^{-1}$ ( $3.27 \pm 0.64 \text{ PgC yr}^{-1}$ )
LNLGOGIS	$19.97 \pm 0.18 \text{ PgCO}_2 \text{ yr}^{-1}$ ( $5.45 \pm 0.05 \text{ PgC yr}^{-1}$ )	$4.03 \pm 2.36 \text{ PgCO}_2 \text{ yr}^{-1}$ ( $1.10 \pm 0.64 \text{ PgC yr}^{-1}$ )	$11.54 \pm 1.79 \text{ PgCO}_2 \text{ yr}^{-1}$ ( $3.15 \pm 0.49 \text{ PgC yr}^{-1}$ )
GCB 2021	$19.8 \pm 0.73 \text{ PgCO}_2 \text{ yr}^{-1}$ ( $5.39 \pm 0.2 \text{ PgC yr}^{-1}$ )	$6.82 \pm 3.15 \text{ PgCO}_2 \text{ yr}^{-1}$ ( $1.86 \pm 0.86 \text{ PgC yr}^{-1}$ )	$10.67 \pm 1.83 \text{ PgCO}_2 \text{ yr}^{-1}$ ( $2.91 \pm 0.5 \text{ PgC yr}^{-1}$ )

$\Delta C_{\text{loss}}$ , suggesting that terrestrial carbon stocks may be decreasing. The LNLG and IS results also differ most in the tropics, with LNLG suggesting greater terrestrial carbon stock loss over  $0^\circ$ – $30^\circ$  N but less over  $0^\circ$ – $30^\circ$  S. The differences in  $\text{CO}_2$  fluxes between these experiments are not well understood, and both experiments evaluate well against independent data (Sect. 4).

The spatial distribution of NCE over 2015–2020 at  $1^\circ \times 1^\circ$  and aggregated to country-scale for the LNLGIS experiment is shown in Fig. 12. At  $1^\circ \times 1^\circ$  (Fig. 12a-b), localized fossil fuel emissions are visible, generally corresponding to urban areas and industrialized regions. These emissions are interspersed over broad source and sink structures that are driven by biosphere removals or emissions. Land biosphere removal is most evident across the northern mid-high latitudes. In contrast, tropical removals and emissions are more regional. When NCE is aggregated to the country-scale (Fig. 12c-d), most countries are net sources driven by fossil fuel emissions, particularly in the northern extratropics. Figure 12e-f shows the 2015–2020 mean country-level  $\Delta C_{\text{loss}}$  for the LNLGIS experiment. Increasing terrestrial carbon stocks (negative  $\Delta C_{\text{loss}}$ ) is found for most extratropical countries, while tropical countries can have gains or losses. Notably, the uncertainty in  $\Delta C_{\text{loss}}$  is larger in the tropics, particularly for mid-sized countries. Overall, small to mid-sized countries generally have uncertainties comparable to the magnitude of  $\Delta C_{\text{loss}}$ , reflecting the fact that atmospheric  $\text{CO}_2$  measurements best constrain fluxes over large scales. Spatial maps of NCE and  $\Delta C_{\text{loss}}$  for each experiment are shown in the supplementary materials (Fig. S4-7).

Differences in NCE and  $\Delta C_{\text{loss}}$  between the v10 OCO-2 MIP experiments can be considerable (the statistical significance of these differences is quantified by the Z statistic, see Sect. 5.1). The underlying cause of differences between the v10 OCO-2 MIP experiments are not well understood, but the differences are likely impacted by the different spatial and temporal distribution of




**Figure 11.** Zonal-mean (a) NCE, (b) FF + lateral fluxes, and (c)  $\Delta C_{\text{loss}}$  for  $30^\circ$  increments of latitude based on  $1^\circ \times 1^\circ$  estimates averaged over 2015–2020. IS, LNLG, LNLGIS and LNLGOGIS median estimates are shown by solid lines and one-sigma uncertainties are shown by the shaded region.

580 LNLG and IS measurements (see Sec. 5.2), model transport errors (Stephens et al., 2007; Schuh et al., 2019, 2022) and residual retrieval biases in the OCO-2  $X_{\text{CO}_2}$  retrievals (Peiro et al., 2022). Unfortunately, the regions showing the largest differences in fluxes generally have few independent atmospheric  $\text{CO}_2$  measurements for validation, limiting our ability to distinguish between different causes. Thus, we believe that NCE and  $\Delta C_{\text{loss}}$  estimates are most reliable when agreement is found across the v10 OCO-2 MIP experiments.

585 We will now show examples of carbon budgets for four countries from this dataset. Figure 13 shows the 2015–2020 mean FF,  $F_{\text{rivers export}}$ ,  $F_{\text{crop trade}}$ ,  $F_{\text{wood trade}}$ ,  $\Delta C_{\text{loss}}$ , and NCE fluxes for the USA, India, Indonesia, and Australia. All of the  $\text{CO}_2$  fluxes on the left of the dashed line combine to give the NCE flux constrained by the v10 OCO-2 MIP experiments. We find that FF is the strongest contributor to NCE for all countries, but that  $\Delta C_{\text{loss}}$  also plays a strong modulating role. For example, negative  $\Delta C_{\text{loss}}$  (increasing terrestrial carbon stocks) for the USA reduces NCE to be less than would be expected given the FF emissions. Conversely, Indonesia has positive  $\Delta C_{\text{loss}}$  (decreasing terrestrial carbon stocks), resulting in increased NCE relative to FF. Some countries also show differences in  $\Delta C_{\text{loss}}$  between v10 OCO-2 MIP experiments. For example, the LNLG and LNLGIS experiments suggest negative  $\Delta C_{\text{loss}}$  for India, while the IS suggest  $\Delta C_{\text{loss}}$  is roughly neutral. Figures of carbon budgets for 28 additional countries (Fig. S8) and 14 regions (Fig. S9) are shown in the supplementary materials.


The carbon budgets can also be examined for individual years (Fig. 14). Both Indonesia and Australia show considerable variations in  $\Delta C_{\text{loss}}$  that drive variations in NCE over this period. Indonesia has a large positive  $\Delta C_{\text{loss}}$  in 2015, driven

595



**Figure 12.** Median ( $\overline{NCE}$ ) and one standard deviation ( $\sigma_{NCE}$ ) of NCE on a (a-b)  $1^\circ \times 1^\circ$  grid and (c-d) aggregated to country-scale for the v10 OCO-2 MIP LNLGIS experiment averaged over 2015–2020. (e-f) Median and one standard deviation of country-scale  $\Delta C_{\text{loss}}$  averaged over 2015–2020 derived from the LNLGIS v10 OCO-2 MIP experiment.

by warm-dry weather and fires during 2015 El Niño (Yin et al., 2016). Australia showed strong negative  $\Delta C_{\text{loss}}$  (except for IS) during 2016, which was the 15th wettest year on record (precipitation 17% above average; Bureau Of Meteorology,



**Figure 13.** CO<sub>2</sub> budget for the USA, India, Indonesia, and Australia averaged over 2015–2020. Bars show the median +/- one standard deviation of FF,  $F_{\text{rivers export}}$  (R),  $F_{\text{crop trade}} + F_{\text{wood trade}}$  (CW),  $\Delta C_{\text{loss}}$ , and NCE (note that these quantities are related through Eqn. 2).


2017). Australia also showed anomalous positive  $\Delta C_{\text{loss}}$  during 2019, which was the warmest and driest year on record, with considerable terrestrial carbon loss related to biomass burning in the southeast (Byrne et al., 2021). Variations in NCE are also found related to FF emissions. In particular, a reduction in NCE is found for 2019 and 2020 in the USA that is primarily linked to a reduction in FF emissions rather than  $\Delta C_{\text{loss}}$ . Timeseries of NCE and  $\Delta C_{\text{loss}}$  for 28 additional countries (Fig. S10, S11) and 14 regions (Fig. S12, S13) are shown in the supplementary materials.

## 8 Discussion

Here we discuss the current limitations of top-down country-level CO<sub>2</sub> budgets and activities that can improve these estimates. Sect. 8.1 discusses current CO<sub>2</sub> observing systems and possible future expansions. Sect. 8.2 discusses current atmospheric CO<sub>2</sub> inversion systems, planned developments, and opportunities for improvement. Finally, Sect. 8.3 discusses remaining challenges in estimating carbon stock changes from atmospheric CO<sub>2</sub> inversions.

### 8.1 Observations

In the context of global inversion analyses, measurements of atmospheric CO<sub>2</sub> best inform annual-mean biosphere–atmosphere CO<sub>2</sub> fluxes over large spatial scales (e.g., continental-to-global) due to rapid mixing in the atmosphere and gaps in current measurement coverage. The confidence in these top-down estimates decreases as we move to smaller spatial scales, with the minimum spatial scale being dependent on the density, precision and sensitivity of the measurements. Future refinements in top-down CO<sub>2</sub> budgets will depend on increasing observational density (Sect. 8.1.1), improved validation (Sect. 8.1.2), and data harmonization (Sect. 8.1.3).



**Figure 14.** Timeseries of the carbon budget for the USA, India, Indonesia, and Australia. Solid lines show the median estimates and shaded areas show +/- one standard deviation.

### 615 8.1.1 Expanding observations

An expanding network of CO<sub>2</sub> observing systems provides an opportunity to reduce uncertainties in top-down estimates of NCE. Across much of the globe, country-level estimates of NCE have been limited by the observational coverage of in situ CO<sub>2</sub> measurements and X<sub>CO<sub>2</sub></sub> retrievals. However, there are a number of planned expansions in observing systems that will help fill data gaps.

620 The first-generation of space-based CO<sub>2</sub> systems currently in operation (GOSAT, GOSAT-2, OCO-2, OCO-3, TanSat) were designed primarily as proof-of-concept missions to demonstrate that space-based measurements could yield X<sub>CO<sub>2</sub></sub> retrievals with the precision and accuracy required to quantify emissions and removals of CO<sub>2</sub>. Planned future missions will expand





and improve upon current observing systems. MicroCarb, a France-UK mission, is expected to start operations in 2023 with an additional spectral band to better characterize the light path for the estimation of  $X_{\text{CO}_2}$  (Bertaux et al., 2020). Japan's  
625 GOSAT-GW mission (<https://gosat-gw.nies.go.jp/en/>), which will be launched in early 2024, will also incorporate improved capabilities for  $\text{CO}_2$  as well as  $\text{CH}_4$ . Soon after, NASA plans to launch the GeoCarb mission (<https://www.ou.edu/geocarb>), which will be hosted on a commercial communications satellite in geostationary orbit at a longitude around  $85^\circ$  W. From that vantage point, GeoCarb can return the data needed to estimate the column average dry air mole fraction of  $\text{CO}_2$ ,  $\text{CH}_4$  and carbon monoxide (CO) over most of North and South America at a spatial resolution of 5 to 10 km every day. In 2025, the  
630 European Copernicus program will begin to deploy the first operational  $\text{CO}_2$  and  $\text{CH}_4$  monitoring constellation, CO2M (Pinty et al., 2017; Janssens-Maenhout et al., 2020). The CO2M constellation will eventually include up to three satellites, flying in formation to collect measurements at 2 km by 2 km resolution over the entire globe at weekly intervals. In addition, a follow-on to the Chinese TanSat mission is currently under development (Yang et al., 2018).

Most current and planned space-based  $\text{CO}_2$  observing systems are passive, in that they rely on reflected sunlight to retrieve  
635  $X_{\text{CO}_2}$ . Active satellite missions, which use lidars for their light source, could provide coverage when reflected sunlight is not available or of insufficient intensity, such as at night and at high latitudes in the winter hemisphere when solar zenith angles are large. These systems also have the potential to better characterize systematic errors in current passive instruments by using pulse timing information to get a better estimate of path length and to filter out scattered light from clouds and aerosols (Abshire et al., 2010).

640 As space-based  $\text{CO}_2$  observing systems expand, sub-orbital discrete air sampling (i.e., flask) and continuous  $\text{CO}_2$  observing systems will remain critical for developing top-down  $\text{CO}_2$  budgets. These in situ observations are the global standard for GHG measurements, because they can undergo direct calibration relative to the WMO  $\text{CO}_2$ -in-air mole fraction scale, which is SI-traceable (Hall et al., 2021). In contrast, open-path remote sensing measurements (both TCCON and satellite) can not be calibrated using standard gasses; they can only be compared to in situ vertical profile observations made relative to the WMO  
645 scale, with the differences used to adjust the remote sensing observations (e.g., Wunch et al., 2011). As such, in situ data are critical for linking remote sensing observations of  $\text{CO}_2$  to the accepted trace gas scales. In situ data also provide complementary observational coverage to space-based observing systems (Byrne et al., 2017). Space-based measurements have broad spatial coverage but with seasonal variations driven by sunlight, and have data gaps in persistently cloudy regions. In contrast, flask and in situ data can be deployed year-round and regardless of cloud cover. Additionally, in situ observations most typically  
650 represent the planetary boundary layer where flux signals in atmospheric  $\text{CO}_2$  are larger than the signal as expressed in the column mean (Feng et al., 2019). Thus, these data play a critical role for improving carbon cycle constraints, especially in high latitude and persistently cloudy regions (such as the tropics), and we encourage an expansion of these systems in these undersampled regions. Regular measurements of  $\text{CO}_2$  using light aircraft above several sites in Amazonia exist (e.g., Gatti et al., 2021; Miller et al., 2021), but these measurement records, as well as a nascent aircraft program in Uganda, have been so  
655 far funded using short-term grants.

Measurements of stable- ( $^{13}\text{C}/^{12}\text{C}$ ) and radio- ( $^{14}\text{C}/\text{C}$ ) isotope ratios of carbon in  $\text{CO}_2$  provide powerful tools for source attribution. Radiocarbon is absent from fossil fuels making it ideal for distinguishing fossil versus biologic carbon fluxes, and



inversions using measurements of CO<sub>2</sub> and <sup>14</sup>C/C have been used to provide top-down constraints on national-scale fossil CO<sub>2</sub> emissions (Basu et al., 2020). Atmospheric <sup>13</sup>C/<sup>12</sup>C ratios provide insight into ecosystem stress and its relation to climate via  
660 constraint of ecosystem water use efficiency (photosynthesis relative to water loss by transpiration) and has been used in box models (Keeling et al., 2017) and inversions (Peters et al., 2018). Atmospheric <sup>13</sup>C/<sup>12</sup>C ratio data are generally available where discrete air samples are collected by various networks, but <sup>14</sup>C/C ratio data are more limited as they tend to require larger samples and measurement costs are greater. Other tracers closely related to CO<sub>2</sub>, such as O<sub>2</sub>/N<sub>2</sub> (Keeling and Graven, 2021) and Carbonyl Sulphide (e.g., Hu et al., 2021; Remaud et al., 2022) are also limited yet provide valuable information on global  
665 ocean/NBE and regional-scale photosynthesis/respiration partitioning, respectively. Increasing the temporal and spatial density of these data, particularly across poorly sampled regions, will allow for more diagnostic power of carbon cycle processes than is possible with CO<sub>2</sub> alone.

### 8.1.2 Data validation

Validation of X<sub>CO<sub>2</sub></sub> retrievals is critical for ensuring that retrieval biases do not strongly impact flux estimates. Current gaps in  
670 coverage of ground-based and airborne measurements have limited our confidence in flux inferences from space-based data. For example, large CO<sub>2</sub> emissions over northern Sub-Saharan Africa are a robust feature of the inversions that assimilate satellite X<sub>CO<sub>2</sub></sub> retrievals (Palmer et al., 2019), but there are few independent CO<sub>2</sub> measurements to confirm whether this inference is a real signal or an artifact of regional retrieval biases. Increased validation of space-based observations will also provide critical support for improved space-based inferences. Space-based measurements rely on validation against ground  
675 based X<sub>CO<sub>2</sub></sub> retrievals from the TCCON (Wunch et al., 2011) and the COllaborative Carbon Column Observing Network (COCCON, Frey et al., 2019). In turn, these sites rely on in situ CO<sub>2</sub> measurements from aircraft profiles and AirCore (Karion et al., 2010) to tie their measurements to the WMO scale (Wunch et al., 2010; Messerschmidt et al., 2011). These data have been critical for validating and improving X<sub>CO<sub>2</sub></sub> retrievals (Wunch et al., 2017b; O'Dell et al., 2018; Kiel et al., 2019). Continued funding of these activities will be crucial for improving top-down CO<sub>2</sub> flux estimates and expansion of these observing systems  
680 into undersampled regions, such as the tropics and high latitudes, will also be important for identifying and addressing residual X<sub>CO<sub>2</sub></sub> retrieval biases. In addition, efforts to cross-calibrate TCCON and COCCON sites will be helpful for minimizing site-to-site biases and identifying spurious drifts in X<sub>CO<sub>2</sub></sub>. We encourage future campaigns aimed at site-to-site comparisons similar to the FRM4GHG campaign that deployed total column GHG traveling standard instruments at several TCCON sites as part of ESA's FRM4GHG-2 project (Sha et al., 2020).

### 8.1.3 Data harmonization

Further advancements in top-down flux estimates will be possible through combining the observational constraints from the constellation space-based sensors and ground-based instruments. Assimilating these data concurrently within inversion systems will increase our ability to recover net fluxes over smaller regions. However, these instruments must be cross-calibrated against common standards to use these data together, as small inter-calibration differences could potentially strongly impact flux  
690 estimates. We encourage support of these critical cross-calibration activities, as are outlined in Crisp et al. (2018).



## 8.2 Atmospheric CO<sub>2</sub> inversions

Atmospheric CO<sub>2</sub> inversion analyses are a critical tool for estimating surface fluxes from observations of atmospheric CO<sub>2</sub>. Expanding observational coverage provides both opportunities and challenges for inversion systems. By addressing the current limitations of our inversion systems, we will be able to take full advantage of increasing observations to improve country-level top-down estimates of NCE and  $\Delta C_{\text{loss}}$ . Here we discuss ongoing and planned developments (Sect. 8.2.1), improving model transport (Sect. 8.2.2), missing processes and required assumptions (Sect. 8.2.3), and uncertainty quantification (Sect. 8.2.4).

### 8.2.1 Ongoing and planned developments

To date, there are four operational or quasi-operational atmospheric CO<sub>2</sub> inversion systems: CarbonTracker (Jacobson et al., 2020), CAMS (Chevallier et al., 2005b), Jena CarboScope (Rödenbeck et al., 2018) and CMS-Flux (Liu et al., 2021a) that are regularly updated on annual or quarterly timescales. These systems produce NBE and air-sea flux estimates from either in situ CO<sub>2</sub> measurements (CarbonTracker, Jena CarboScope), OCO-2 X<sub>CO<sub>2</sub></sub> retrievals (CMS-Flux) or both (CAMS). Similarly, there are seven inversion models (including the aforementioned models) that update CO<sub>2</sub> flux estimates annually for the Global Carbon Budget (Friedlingstein et al., 2022), including CAMS (Chevallier et al., 2005b), CarbonTracker Europe (CTE van der Laan-Luijkx et al., 2017), Jena CarboScope (Rödenbeck et al., 2018), UoE in situ (Feng et al., 2016), NISMON-CO2 (Niwa et al., 2017), MIROC4-ACTM (Saeki and Patra, 2017; Chandra et al., 2021), and CMS-Flux (Liu et al., 2021a).

The OCO-2 MIP activities have semi-regularly performed ensemble inversion experiments (Crowell et al., 2019; Peiro et al., 2022). To date, OCO-2 MIP experiments have been linked to new versions of the ACOS retrieval algorithm, with major improvements to the quality of X<sub>CO<sub>2</sub></sub> retrievals occurring during each update. However, as the quality of retrievals have improved (particularly for ACOS v10 onwards), updates to the ACOS retrieval algorithm are becoming less of a driver for new OCO-2 MIP experiments. In the future, OCO-2 MIP activities could become more regular with annual updates.

The first top-down CO<sub>2</sub> system for use in inventory development is CarbonWatch-NZ, under development in New Zealand (<https://niwa.co.nz/climate/research-projects/carbon-watch-nz>). This program includes expanded CO<sub>2</sub> measurement sites and the development of a regional atmospheric CO<sub>2</sub> inverse system to quantify the carbon budgets of New Zealand's forest, grassland and urban environments. Initial results suggest stronger uptake by intact forests than estimated through bottom-up estimates (Steinkamp et al., 2017). This system may serve as an example for other nations through the Integrated Global Greenhouse Gas Information System (IG3IS) framework.

Beyond existing activities, there are a number of planned projects. The European Commission's Copernicus program (<https://www.copernicus.eu>) has a number of developments ongoing and planned, particularly in building anthropogenic CO<sub>2</sub> emissions monitoring and verification support capacity (CO2MVS; Janssens-Maenhout et al., 2020), which is directly linked to the development and launch of the new CO2M mission and is expected to be operational from 2026 onwards. Further, there are a number of recently completed, ongoing, and planned projects to develop and improve inversion systems to develop operational capacity. Examples include the recently completed CO<sub>2</sub> Human Emissions (CHE) project (<https://www.che-project.eu/>) and follow-up CoCO2 project (<https://coco2-project.eu/>) that is ongoing, as-well as the VERIFY project (<https://verify.lsce.ipsl.fr/>).



725 These projects are developing and refining inversion systems to estimate anthropogenic fossil fuel emissions as well as emis-  
sions and removals from the AFOLU sector. Future planned projects include developing approaches to utilize co-emitted  
species and auxiliary observations ( $^{14}\text{C}$ , solar induced fluorescence, Carbonyl Sulfide, and others) in order to isolate some of  
the  $\text{CO}_2$  budget components and improve our understanding of the carbon cycle. For example, multiple data streams could be  
used together to optimize the dynamic global vegetation model parameters (e.g., Peylin et al., 2016).

730 In contrast to recent European efforts, there is no mandate for an operational top-down carbon flux-attribution system  
in the US. Nevertheless, efforts at NOAA centered around CarbonTracker (Jacobson et al., 2020) have been able to pro-  
duce NBE estimates with relatively low latency harnessing the Agency's substantial flask and in situ  $\text{CO}_2$  network. In ad-  
dition, NOAA has developed a higher spatial resolution North American regional inverse system, CarbonTracker-Lagrange  
(<https://gml.noaa.gov/ccgg/carbontracker-lagrange/>; Hu et al., 2019). In anticipation of the launch of OCO in 2009, NASA has  
supported research and development efforts needed to prototype an operational flux estimation system. In particular, the Carbon  
735 Monitoring System program (<https://carbon.nasa.gov/>) has led to the development of both low-latency (2 month) atmospheric  
 $\text{CO}_2$  reanalysis (Weir et al., 2021) and approaches to combine top-down NCE estimates with other trace gas measurements  
(e.g., CO) and non-atmospheric carbon data (e.g., above-ground biomass) to provide improved understanding of carbon cycle  
processes (Liu et al., 2017; Byrne et al., 2020, 2021; Bloom et al., 2020). There is substantial technical capacity to build an  
operational system but requires a coordinated effort between federal agencies, academia, and private interests.

740 In Canada, a prototype operational regional inverse modeling system, the Environment and Climate Change Canada (ECCC)  
National Carbon Flux Inversion System (ENCIS), is being developed to provide quantitative information on  $\text{CO}_2$  (and  $\text{CH}_4$ )  
flux estimates over Canada from national to provincial scales, as well as to understand the carbon cycle in Canada such as  $\text{CO}_2$   
flux in boreal managed and unmanaged forests, wetland emissions of  $\text{CH}_4$ , and GHG emissions over a potentially thawing  
permafrost in response to the climate change. ENCIS is a regional inverse modeling system based on Lagrangian approach and  
745 driven by metrology from the Global Environmental Multiscale (GEM) model (Girard et al., 2014) and is expected to have  
 $1^\circ \times 1^\circ$  spatial resolution.

### 8.2.2 Improving CTM transport

Errors in the representation of atmospheric transport by CTMs has long been recognized as a major source of error in atmo-  
sphere  $\text{CO}_2$  inversion analyses (Law et al., 1996; Law and Simmonds, 1996; Denning et al., 1995, 1999a, b; Baker et al.,  
750 2006a; Stephens et al., 2007). Improvements to model transport will provide critical improvements to NCE and  $\Delta C_{\text{loss}}$  es-  
timates. Systematic errors in model transport limit our ability to relate surface fluxes and  $\text{CO}_2$  observations, and can lead to  
incorrect inferences of surface fluxes (Yu et al., 2018; Schuh et al., 2019; Stanevich et al., 2020). Improving model transport  
will require work in two areas: (1) improving model parameterizations of unresolved transport, particularly in coarse off-  
line CTMs (like GEOS-Chem run at  $4^\circ \times 5^\circ$  in this ensemble) where the spatial and temporal coarsening of meteorological  
755 fields can “average-out” vertical transport that is resolved in the parent model (Yu et al., 2018; Stanevich et al., 2020); and  
(2) increasing spatial and temporal resolution in model simulations, which can better resolve atmospheric transport processes  
(Agustí-Panareda et al., 2019; Schuh et al., 2019). However, it should be noted that there are limitations to the improvements



that can come from increased model resolution in the global inversion context due to underlying meteorological uncertainties (Liu et al., 2011; Polavarapu et al., 2016, 2018; McNorton et al., 2020). Computational cost is also a significant challenge in  
760 inversion systems, because transport models usually scale poorly on supercomputers, for example because of the volume of meteorological data required as input.

As transport models are refined, it will be critical to periodically test their ability to represent large scale atmospheric dynamics. This can be tested using long-lived trace gas species, including sulfur hexafluoride (Schuh et al., 2019), idealized age of air tracer (Krol et al., 2018), and beryllium-7 (Stanevich et al., 2020). Simulations of these trace species are critical in  
765 the context of inversion MIPs to gauge inter-model variability and average model bias (Schuh et al., 2019). Similarly, Rn 222 is a useful short lived gas species that enables modelers to evaluate the vertical mixing within the column (Remaud et al., 2018). In addition, model intercomparison studies have proven useful for diagnosing transport errors (e.g., Gaubert et al., 2019; Zhang et al., 2022), and we recommend further activities, such as within the Atmospheric Tracer Transport Model Intercomparison Project (TRANSCOM) framework.

### 770 8.2.3 Missing processes and required assumptions

The flux estimates provided here do not explicitly account for the atmospheric-chemical production of atmospheric CO<sub>2</sub>, which occurs from the oxidation of reduced carbon gasses. Instead, these fluxes are either prescribed as surface-atmosphere fluxes (e.g., for FF CO emissions) or neglected from the prior fluxes. This can cause inverse modeling systems to implicitly incorporate the atmospheric CO<sub>2</sub> source in optimized surface-atmosphere emissions and removals (i.e. air-sea fluxes and  
775 NBE), which can be far from the actual source of the reduced gas. For example, FF CO emissions are largely emitted in the northern extratropics but largely oxidized to CO<sub>2</sub> in the tropical troposphere. These incorrectly located emissions of CO<sub>2</sub> are large enough to impact top-down inversions (Enting and Mansbridge, 1991; Suntharalingam et al., 2005; Nassar et al., 2010; Wang et al., 2020). Future studies that aim to incorporate an atmospheric source of CO<sub>2</sub> would help correct for this current spatial bias (Ciais et al., 2022).

780 A critical assumption in the top-down CO<sub>2</sub> budgets estimated here has been that FF emissions are known and unbiased. Uncertainties in inventory-based FF emission estimates at global and country levels (e.g., Andres et al., 2014) are smaller than top-down NCE estimates; however, inventory-based emission estimates are prone to systematic biases due to the nature of the estimation approach (Guan et al., 2012; Oda et al., 2019) and FF uncertainties could bias the partitioning of NCE between FF and NBE (and propagate into  $\Delta C_{\text{loss}}$ ) over countries with large emissions and lower reliability of statistical data collection  
785 system, such as China. For example, Saeki and Patra (2017) show that an inferred increase in removals of CO<sub>2</sub> by the biosphere over China during 2001–2010 are likely to be an artifact imposed by an error in the trend of anthropogenic CO<sub>2</sub> emissions.

### 8.2.4 Uncertainty quantification

The uncertainty in NCE reported here is an estimate of the standard deviation of the v10 OCO-2 MIP ensemble members. This is meant to characterize uncertainties originating from the inversion configuration (such as the transport model, inver-  
790 sion method, and prior constraints). However, there are also limitations to this method. First, there is only a small ensemble



of 11 MIP ensemble members included in this analysis, and an over-representation of inversions using two transport models: TM5 (3) and GEOS-Chem (5), which makes uncertainty quantification challenging. Future approaches that employ “borrowing strength” (Mearns and et al., 2007; Cressie and Kang, 2016) could be employed to better characterize ensemble uncertainty. Second, the ensemble-based uncertainty does not capture some sources of uncertainty. In particular, Bayesian posterior uncertainties are not considered here (see Sect. 5.2), due to the fact that many of the inversion systems participating in the v10 OCO-2 MIP do not calculate this uncertainty. In addition, we find that the ensemble members that produce Bayesian uncertainties show large differences in magnitude. Thus, this is an area of future improvement for MIP activities, and we recommend more work into characterizing this error component in ensemble inversion experiments. We also note that using an analytic framework, posterior uncertainties and their sensitivities to prior information could be further examined, as has been done for methane (Worden et al., 2022).

### 8.3 Stockchange estimates

AFOLU emissions and removals are generally quantified as terrestrial carbon stockchanges in managed lands. A number of challenges remain in estimating this quantity from top-down methods. Firstly, lateral fluxes of carbon remain quite uncertain (and associated uncertainty estimates are themselves quite uncertain). The best constrained lateral fluxes are annual country-level  $F_{\text{wood trade}}$  and  $F_{\text{wood trade}}$ , which are reported to the UN Food and Agriculture Organization. These fluxes are more uncertain on sub-national scales and sub-annual timescales. Meanwhile,  $F_{\text{rivers export}}$  is best quantified on basin scales, where stream gauge measurements inform carbon fluxes. Improving sub-national and sub-annual estimates of lateral fluxes would have several benefits: first, this would allow for better sub-national attribution, where regional fluxes could be better quantified. Second, this would allow for incorporating the atmospheric imprint of these carbon fluxes as prior information within atmospheric CO<sub>2</sub> inversion analyses, which may improve flux estimates on sub-national scales.

The Global Stocktake and Paris Agreement do not consider emissions and removals from unmanaged lands. Separating managed lands from unmanaged lands is top-down NCE remains a major challenge, given the smoothed large-scale CO<sub>2</sub> flux constraints provided by these top-down methods and the fact that both managed and unmanaged lands can experience considerable stock changes driven by interannual climate variations (e.g., El Niño) and in response to rising CO<sub>2</sub> and climate change. In addition, separating managed and unmanaged lands is further complicated by the fact that there is considerable ambiguity in the definitions managed lands, which can also vary by country (Grassi et al., 2018; Chevallier, 2021). We recommend that each party provide a mask to unambiguously define the plots considered as managed from year to year (Chevallier, 2021).

## 9 Conclusions

We introduced a pilot top-down CO<sub>2</sub> budget dataset (Byrne et al., 2022) intended to start a dialogue between research communities and to identify ways that top-down flux estimates can inform country-level carbon budgets. This dataset provides annual country-level and 1° × 1° gridded top-down NCE and  $\Delta C_{\text{loss}}$  over 2015–2020, in addition to bottom-up FF and lateral fluxes. These data are provided for four experiments from the v10 OCO-2 MIP that differ in the data used in the assimilation:



IS, LNLG, LNLGIS, and LNLGOGIS. In addition, we provide two metrics for interpreting country-level estimates: (1) the Z-statistic (Sect. 5.1), which quantifies the agreement between IS and LNLG NCE estimates, and (2) the FUR (Sect. 5.2), which quantifies the impact of atmospheric CO<sub>2</sub> data in reducing flux uncertainties.

Country-level flux estimates generally show robust signals for large extratropical countries (e.g., USA, Russia, China). Agreement between the experiments generally decreases for mid-sized countries (e.g., Turkey), particularly in regions with sparse observational coverage for the in situ network (such as the tropics). Large divergences between the IS and LNLG experiments occur in some regions, particularly northern Sub-Saharan Africa, and could be related to the sparsity of in situ CO<sub>2</sub> measurements or biases in OCO-2 retrievals. However, the sparsity of independent CO<sub>2</sub> measurements in these regions precludes definitive conclusions. We urge caution in interpreting the 1° × 1° gridded results and suggest collaborating with experts in atmospheric CO<sub>2</sub> inversion systems when using those data.

The accuracy of top-down NCE estimates were characterized through comparisons against independent atmospheric CO<sub>2</sub> datasets, and through comparisons against pCO<sub>2</sub>-based air–sea CO<sub>2</sub> fluxes. Overall, the IS, LNLG, and LNLGIS were found to show the best agreement against independent CO<sub>2</sub> measurements, and we recommend using these experiments for analysis. Poorer agreement for experiments assimilating OCO-2 ocean glint X<sub>CO<sub>2</sub></sub> retrievals, suggesting that residual retrieval biases adversely impact the LNLGOGIS experiment and we urge caution in interpreting these data.

For future GSTs, top-down NCE estimates will be refined as new space-based X<sub>CO<sub>2</sub></sub> observing systems expand and retrieval algorithms are improved. Complementary expansions of ground-based and aircraft-based CO<sub>2</sub> measurements in under-sampled regions will similarly fill critical observational gaps in regions with large uncertainties and susceptibility to retrieval biases. Improvements to atmospheric CO<sub>2</sub> inversion systems, including reductions to systematic transport errors and improved error characterization, will be critical for refining top-down CO<sub>2</sub> budgets. And improved estimates of lateral carbon fluxes and managed lands maps will refine estimates of AFOLU emissions and removals.

## 10 Data availability

Top-down CO<sub>2</sub> budgets (Byrne et al., 2022) are available from the Committee on Earth Observation Satellites' (CEOS) website: <https://doi.org/10.48588/npf6-sw92>. Gridded NBE and air-sea fluxes from the OCO-2 MIP are available at [https://gml.noaa.gov/ccgg/OCO2\\_v10mip/](https://gml.noaa.gov/ccgg/OCO2_v10mip/). Fossil fuel emissions prescribed in the inversions can be downloaded from <https://zenodo.org/record/4776925#.YNX96hNKj2U>. The ODIAC2020 emission data product can be downloaded from the Global Environmental Database hosted by the Center for Global Global Environmental Research at NIES ([https://db.cger.nies.go.jp/dataset/ODIAC/DL\\_odiad2020.html](https://db.cger.nies.go.jp/dataset/ODIAC/DL_odiad2020.html)). SeaFlux pCO<sub>2</sub>-based air–sea fluxes were downloaded from <https://zenodo.org/record/5482547#.Yowg18ZID1I>, accessed 23 May 2022.



## Appendix A: TCCON sites

**Table A1.** TCCON sites used for evaluation of posterior CO<sub>2</sub> fields of the v10 OCO-2 MIP experiments.

TCCON site	Country	Latitude	Longitude	Reference
Eureka	Canada	80.05° N	86.42 °W	Strong et al. (2019)
Ny-Ålesund	Norway	78.9° N	11.9 °E	Notholt et al. (2019b)
Sodankylä	Finland	67.4° N	26.6 °E	Kivi et al. (2014)
East Trout Lake	Canada	54.4° N	105.0 °W	Wunch et al. (2017a)
Bremen	Germany	53.10° N	8.85 °E	Notholt et al. (2019a)
Karlsruhe	Germany	49.1° N	8.4 °E	Hase et al. (2014)
Paris	France	48.8° N	2.4 °E	Te et al. (2014)
Orléans	France	47.9° N	2.1 °E	Warneke et al. (2019)
Garmisch	Germany	47.5° N	11.1 °E	Sussmann and Rettinger (2018a)
Zugspitze	Germany	47.3° N	11.0 °E	Sussmann and Rettinger (2018b)
Park Falls	USA	45.9° N	90.3 °W	Wennberg et al. (2017)
Rikubetsu	Japan	43.5° N	143.8 °E	Morino et al. (2014)
Lamont	USA	36.6° N	97.5 °W	Wennberg et al. (2016b)
Anmeiyondo	Korea	36.5° N	126.3 °E	Goo et al. (2014)
Tsukuba	Japan	36.1° N	140.1 °E	Morino et al. (2018a)
Nicosia	Cyprus	35.1° N	33.4 °E	Petri et al. (2020)
Edwards	USA	34.2° N	118.2 °W	Iraci et al. (2016)
JPL	USA	34.2° N	118.2 °W	Wennberg et al. (2016a)
Caltech	USA	34.1° N	118.1 °W	Wennberg et al. (2014)
Saga	Japan	33.2° N	130.3 °E	Kawakami et al. (2014)
Hefei	China	31.9° N	117.2 °E	Liu et al. (2018)
Izaña	Spain	28.3° N	16.5 °W	Blumenstock et al. (2017)
Burgos	Philippines	18.5° N	120.7 °E	Morino et al. (2018b)
Manaus	Brazil	3.2° N	60.6 °W	Dubey et al. (2014)
Ascension Island	UK	7.9° S	14.3 °W	Feist et al. (2014)
Darwin	Australia	12.4° S	130.9 °E	Griffith et al. (2014a);
Réunion island	France	20.9° S	55.5 °W	De Mazière et al. (2017)
Wollongong	Australia	34.4° S	150.9 °E	Griffith et al. (2014b)
Lauder 125HR	New Zealand	45.0° S	169.7 °E	Sherlock et al. (2014)





*Author contributions.* The study was conceived of by DC, and designed by BB, DB, SB, KWB, AC, FC, PC, NC, DC, LH, ARJ, JL, JBM, TO, PKP, BP, AS, CS, and JRW. The v10 OCO-2 MIP experiments were performed by DFB, SB, MB, FC, NC, SC, FD, ARJ, RJ, MSJ, DBAJ, JL, ZL, SM, SMM, RM, AS, AZM, and NZ. Lateral flux estimates were performed by FC, PC, ZD, HT, and YY. FF emissions estimates were performed by TO and SB. TCCON data were collected by NMD, MKD, OEG, BH, RK, IM, JN, YSO, HO, CP, KS, KS, YT, VAV, MV, TW and DW. Evaluation of v10 OCO-2 MIP against co-samples was performed by BB, SB, AC, and ARJ. BB wrote the paper and prepared the figures, with contributions from all co-authors.

*Competing interests.* no competing interests are present

*Acknowledgements.* Part of this research was carried out at the Jet Propulsion Laboratory, California Institute of Technology, under a contract with the National Aeronautics and Space Administration (80NM0018D004). The v10 OCO-2 MIP activity was supported by the NASA OCO Science Team program. BB and JL were supported by the NASA OCO2/3 science team program NNX17ZDA001N-OCO2. ARJ, AS, and DFB were funded by NASA award 80NSSC21K1080. AS was also supported by the NASA grant NNX15AG93G. SC acknowledge from the NASA OCO Science Team grant 80NSSC21K1077. The research of NC, AZM, and MB was supported by Australian Research Council Discovery Project DP190100180 and by NASA ROSES grant 20-OCOST20-0004. AZM is also supported by the Australian Research Council Discovery Early Career Research Award DE180100203. Contribution of AC was supported by NASA ROSES Grant numbers 80NSSC20K0006 and 80NSSC21K1068. MSJ acknowledges the internal funding from NASA's Earth Science Research and Analysis Program. The contributions of FC and MR were supported by the Copernicus Atmosphere Monitoring Service, implemented by the European Centre for Medium-Range Weather Forecasts on behalf of the European Commission (grant no. CAMS73), and by the European Union's Horizon 2020 research and innovation programme under grant agreement No 958927 (Prototype system for a Copernicus CO2 service). It was granted access to the HPC resources of TGCC under the allocation A0110102201 made by GENCI. SF at PNNL is supported by the NASA Carbon Monitoring Program (Grant number: 80HQTR21T0069). The Pacific Northwest National Laboratory is operated by Battelle Memorial Institute under contract DE-AC05-76RL01830. NZ acknowledges support from NOAA (NA18OAR4310266) and NASA (80NSSC18K0908). PKP is partly supported by Environment Research and Technology Development Fund (JPMEERF21S20800) of the Environmental Restoration and Conservation Agency of Japan. SMM was supported by the NASA grants 80NSSC18K0976 and 80NSSC21K1073. NMD was funded by ARC Future Fellowship FT180100327

The TCCON Nicosia site has received additional support from the European Union's Horizon 2020 research and innovation programme (grant agreement No. 856612 /EMME-CARE), the Cyprus Government, and the University of Bremen. The TCCON Anmyeondo site has been funded by the Korea Meteorological Administration Research and Development Program "Developing Technology for Integrated Climate Change Monitoring and Analysis" under grant(KMA2018-00320). TCCON sites at Tsukuba, Rikubetsu and Burgos are supported in part by the GOSAT series project. Burgos is supported in part by the Energy Development Corp. Philippines. The Eureka TCCON measurements were made at the Polar Environment Atmospheric Research Laboratory (PEARL) by the Canadian Network for the Detection of Atmospheric Change (CANDAC), primarily supported by the Natural Sciences and Engineering Research Council of Canada, Environment and Climate Change Canada, and the Canadian Space Agency. The TCCON site at Réunion Island has been operated by the Royal Belgian Institute for Space Aeronomy with financial support since 2014 by the EU project ICOS-Inwire and the ministerial decree for ICOS (FR/35/IC1 to FR/35/C6) and local activities supported by LACy/UMR8105 and by OSU-R/UMS3365 – Université de La Réunion. Darwin



and Wollongong TCCON stations are supported by ARC grants DP160100598, LE0668470, DP140101552, DP110103118 and DP0879468 and NASA grants NAG512247 and NNG05GD07G.

890 We thank Robert J. Andres for providing uncertainty estimates for CDIAC fossil fuel emission estimates. We thank the data providers of the SeaFlux ensemble for making their pCO<sub>2</sub>-based air–sea CO<sub>2</sub> fluxes publicly available. We are grateful for the leadership of Annemarie Eldering and Mike Gunson of the OCO-2 mission, whose hard work has made this dataset possible.



## References

- Abshire, J. B., Riris, H., Allan, G. R., Weaver, C. J., Mao, J., Sun, X., Hasselbrack, W. E., Kawa, S. R., and Biraud, S.: Pulsed airborne lidar measurements of atmospheric CO<sub>2</sub> column absorption, *Tellus B*, 62, 770–783, <https://doi.org/10.1111/j.1600-0889.2010.00502.x>, 2010.
- Agustí-Panareda, A., Diamantakis, M., Massart, S., Chevallier, F., Muñoz Sabater, J., Barré, J., Curcoll, R., Engelen, R., Langerock, B., Law, R. M., Loh, Z., Morguí, J. A., Parrington, M., Peuch, V.-H., Ramonet, M., Roehl, C., Vermeulen, A. T., Warneke, T., and Wunch, D.: Modelling CO<sub>2</sub> weather – why horizontal resolution matters, *Atmos. Chem. Phys.*, 19, 7347–7376, <https://doi.org/10.5194/acp-19-7347-2019>, 2019.
- Anav, A., Friedlingstein, P., Beer, C., Ciais, P., Harper, A., Jones, C., Murray-Tortarolo, G., Papale, D., Parazoo, N. C., Peylin, P., Piao, S., Sitch, S., Nicolas, V., Andy, W., and Zhao, M.: Spatiotemporal patterns of terrestrial gross primary production: A review, *Rev. Geophys.*, 53, 785–818, <https://doi.org/10.1002/2015RG000483>, 2015.
- Andres, R. J., Boden, T. A., and Higdon, D.: A new evaluation of the uncertainty associated with CDIAC estimates of fossil fuel carbon dioxide emission, *Tellus B*, 66, 23 616, <https://doi.org/10.3402/tellusb.v66.23616>, 2014.
- Baier, B., Sweeney, C., Tans, P., Newberger, T., Higgs, J., and Wolter, S.: NOAA AirCore atmospheric sampling system profiles (Version 20210813), <https://doi.org/https://doi.org/10.15138/6AV0-MY81>, 2021.
- Baker, D., Law, R. M., Gurney, K. R., Rayner, P., Peylin, P., Denning, A., Bousquet, P., Bruhwiler, L., Chen, Y.-H., Ciais, P., et al.: TransCom 3 inversion intercomparison: Impact of transport model errors on the interannual variability of regional CO<sub>2</sub> fluxes, 1988–2003, *Global Biogeochem. Cy.*, 20, <https://doi.org/10.1029/2004GB002439>, 2006a.
- Baker, D., Bösch, H., Doney, S., O'Brien, D., and Schimel, D.: Carbon source/sink information provided by column CO<sub>2</sub> measurements from the Orbiting Carbon Observatory, *Atmos. Chem. Phys.*, 10, 4145–4165, <https://doi.org/10.5194/acp-10-4145-2010>, 2010.
- Baker, D. F., Doney, S. C., and Schimel, D. S.: Variational data assimilation for atmospheric CO<sub>2</sub>, *Tellus B*, 58, 359–365, <https://doi.org/10.5194/acp-10-4145-2010>, 2006b.
- Baker, D. F., Bell, E., Davis, K. J., Campbell, J. F., Lin, B., and Dobler, J.: A new exponentially decaying error correlation model for assimilating OCO-2 column-average CO<sub>2</sub> data using a length scale computed from airborne lidar measurements, *Geosci. Model Dev.*, 15, 649–668, <https://doi.org/10.5194/gmd-15-649-2022>, 2022.
- Ballantyne, A. P., Alden, C. B., Miller, J. B., Tans, P. P., and White, J.: Increase in observed net carbon dioxide uptake by land and oceans during the past 50 years, *Nature*, 488, 70–72, <https://doi.org/10.1038/nature11299>, 2012.
- Basu, S., Lehman, S. J., Miller, J. B., Andrews, A. E., Sweeney, C., Gurney, K. R., Xu, X., Southon, J., and Tans, P. P.: Estimating US fossil fuel CO<sub>2</sub> emissions from measurements of <sup>14</sup>C in atmospheric CO<sub>2</sub>, *Proceedings of the National Academy of Sciences*, 117, 13 300–13 307, <https://doi.org/10.1073/pnas.1919032117>, 2020.
- Bertaux, J.-L., Hauchecorne, A., Lefèvre, F., Bréon, F.-M., Blanot, L., Jougllet, D., Lafrique, P., and Akaev, P.: The use of the 1.27 μm O<sub>2</sub> absorption band for greenhouse gas monitoring from space and application to MicroCarb, *Atmos. Meas. Tech.*, 13, 3329–3374, <https://doi.org/10.5194/amt-13-3329-2020>, 2020.
- Bloom, A. A., Bowman, K. W., Liu, J., Konings, A. G., Worden, J. R., Parazoo, N. C., Meyer, V., Reager, J. T., Worden, H. M., Jiang, Z., Quetin, G. R., Smallman, T. L., Exbrayat, J.-F., Yin, Y., Saatchi, S. S., Williams, M., and Schimel, D. S.: Lagged effects regulate the inter-annual variability of the tropical carbon balance, *Biogeosciences*, 17, 6393–6422, <https://doi.org/10.5194/bg-17-6393-2020>, 2020.



- Blumenstock, T., Hase, F., Schneider, M., García, O. E., and Sepúlveda, E.: TCCON data from Izana (ES), Release GGG2014.R1, <https://doi.org/10.14291/TCCON.GGG2014.IZANA01.R1>, 2017.
- 930 Bolin, B. and Keeling, C.: Large-scale atmospheric mixing as deduced from the seasonal and meridional variations of carbon dioxide, *J. Geophys. Res.*, 68, 3899–3920, <https://doi.org/10.1029/JZ068i013p03899>, 1963.
- Bureau Of Meteorology: Annual climate statement 2016, <http://www.bom.gov.au/climate/current/annual/aus/2016>, accessed: 10 June 2022, 2017.
- Byrne, B., Jones, D. B. A., Strong, K., Zeng, Z.-C., Deng, F., and Liu, J.: Sensitivity of CO<sub>2</sub> Surface Flux Constraints to Observational  
935 Coverage, *J. Geophys. Res.-Atmos*, 112, 6672–6694, <https://doi.org/10.1002/2016JD026164>, 2017.
- Byrne, B., Liu, J., Bloom, A. A., Bowman, K. W., Butterfield, Z., Joiner, J., Keenan, T. F., Keppel-Aleks, G., Parazoo, N. C., and Yin, Y.: Contrasting regional carbon cycle responses to seasonal climate anomalies across the east-west divide of temperate North America, *Global Biogeochem. Cy.*, 34, e2020GB006598, <https://doi.org/10.1029/2020GB006598>, 2020.
- Byrne, B., Liu, J., Lee, M., Yin, Y., Bowman, K. W., Miyazaki, K., Norton, A. J., Joiner, J., Pollard, D. F., Griffith, D. W., Velasco, V. A.,  
940 Deutscher, N. M., Jones, N. B., and Paton-Walsh, C.: The carbon cycle of southeast Australia during 2019–2020: Drought, fires, and subsequent recovery, *AGU Advances*, 2, e2021AV000469, <https://doi.org/10.1029/2021AV000469>, 2021.
- Byrne, B., Baker, D. F., Basu, S., Bertolacci, M., Bowman, K. W., Carroll, D., Chatterjee, A., Chevallier, F., Ciais, P., Cressie, N., Crisp, D., Crowell, S., Deng, F., Deng, Z., Deutscher, N. M., Dubey, M. K., Feng, S., García, O. E., Herkommer, B., Hu, L., Jacobson, A. R., Janardanan, R., Jeong, S., Johnson, M. S., Jones, D. B. A., Kivi, R., Liu, J., Liu, Z., Maksyutov, S., Miller, J. B., Miller, S. M., Morino, I.,  
945 Notholt, J., Oda, T., O'Dell, C. W., Oh, Y.-S., Ohyama, H., Patra, P. K., Peiro, H., Petri, C., Philip, S., Pollard, D. F., Poulter, B., Remaud, M., Schuh, A., Sha, M. K., Shiomi, K., Strong, K., Sweeney, C., Té, Y., Tian, H., Velasco, V. A., Vrekoussis, M., Warneke, T., Worden, J. R., Wunch, D., Yao, Y., Yun, J., Zammit-Mangion, A., and Zeng, N.: Pilot top-down CO<sub>2</sub> Budget constrained by the v10 OCO-2 MIP Version 1, Committee on Earth Observing Satellites, <https://doi.org/10.48588/npf6-sw92>, Version 1.0, 2022.
- Canadell, J. G., Monteiro, P. M., Costa, M. H., Da Cunha, L. C., Cox, P. M., Alexey, V., Henson, S., Ishii, M., Jaccard, S., Koven, C., Lohila, A., Patra, P. K., Piao, S., Rogelj, J., Syampungani, S., Zaehle, S., and Zickfeld, K.: Global Carbon and other Biogeochemical Cycles and Feedbacks, pp. 673–816, Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, <https://doi.org/10.1017/9781009157896.007>, 2021.
- 950 Caspersen, J. P., Pacala, S. W., Jenkins, J. C., Hurtt, G. C., Moorcroft, P. R., and Birdsey, R. A.: Contributions of land-use history to carbon accumulation in US forests, *Science*, 290, 1148–1151, <https://doi.org/10.1126/science.290.5494.114>, 2000.
- 955 Center for International Earth Science Information Network - CIESIN - Columbia University: Gridded Population of the World, Version 4 (GPWv4) National Identifier Grid, Revision 11., <https://sedac.ciesin.columbia.edu/data/set/gpw-v4-national-identifier-grid-rev11>, accessed 3 March 2021, 2018.
- Chandra, N., Patra, P. K., Niwa, Y., Ito, A., Iida, Y., Goto, D., Morimoto, S., Kondo, M., Takigawa, M., Hajima, T., and Watanabe, M.: Estimated regional CO<sub>2</sub> flux and uncertainty based on an ensemble of atmospheric CO<sub>2</sub> inversions, *Atmos. Chem. Phys. Discussions*,  
960 2021, 1–50, <https://doi.org/10.5194/acp-2021-1039>, 2021.
- Chau, T. T. T., Gehlen, M., and Chevallier, F.: A seamless ensemble-based reconstruction of surface ocean pCO<sub>2</sub> and air–sea CO<sub>2</sub> fluxes over the global coastal and open oceans, *Biogeosciences*, 19, 1087–1109, <https://doi.org/10.5194/bg-19-1087-2022>, 2022.
- Chen, Z., Huntzinger, D. N., Liu, J., Piao, S., Wang, X., Sitch, S., Friedlingstein, P., Anthoni, P., Arneth, A., Bastrikov, V., Goll, D. S., Haverd, V., Jain, A. K., Joetzjer, E., Kato, E., Lienert, S., Lombardozi, D. L., McGuire, P. C., Melton, J. R., Nabel, J. E. M. S., Pongratz, J., Poulter, B., Tian, H., Wiltshire, A. J., Zaehle, S., and Miller, S. M.: Five years of variability in the global carbon cycle: comparing an estimate from



- the Orbiting Carbon Observatory-2 and process-based models, *Environ. Res. Lett.*, 16, 054 041, <https://doi.org/10.1088/1748-9326/abfac1>, 2021a.
- Chen, Z., Liu, J., Henze, D. K., Huntzinger, D. N., Wells, K. C., Sitch, S., Friedlingstein, P., Joetzjer, E., Bastrikov, V., Goll, D. S., Haverd, V., Jain, A. K., Kato, E., Lienert, S., Lombardozzi, D. L., McGuire, P. C., Melton, J. R., Nabel, J. E. M. S., Poulter, B., Tian, H., Wiltshire, A. J., Zaehle, S., and Miller, S. M.: Linking global terrestrial CO<sub>2</sub> fluxes and environmental drivers: inferences from the Orbiting Carbon Observatory 2 satellite and terrestrial biospheric models, *Atmos. Chem. Phys.*, 21, 6663–6680, <https://doi.org/10.5194/acp-21-6663-2021>, 2021b.
- Chevallier, F.: Fluxes of Carbon Dioxide From Managed Ecosystems Estimated by National Inventories Compared to Atmospheric Inverse Modeling, *Geophys. Res. Lett.*, 48, e2021GL093 565, <https://doi.org/10.1029/2021GL093565>, 2021.
- 975 Chevallier, F., Engelen, R. J., and Peylin, P.: The contribution of AIRS data to the estimation of CO<sub>2</sub> sources and sinks, *Geophys. Res. Lett.*, 32, <https://doi.org/10.1029/2005GL024229>, 2005a.
- Chevallier, F., Fisher, M., Peylin, P., Serrar, S., Bousquet, P., Bréon, F.-M., Chédin, A., and Ciais, P.: Inferring CO<sub>2</sub> sources and sinks from satellite observations: Method and application to TOVS data, *J. Geophys. Res.-Atmos.*, 110, <https://doi.org/10.1029/2005JD006390>, 2005b.
- 980 Chevallier, F., Breon, F.-M., and Rayner, P. J.: Contribution of the Orbiting Carbon Observatory to the estimation of CO<sub>2</sub> sources and sinks: Theoretical study in a variational data assimilation framework, *J. Geophys. Res.*, 112, <https://doi.org/10.1029/2006JD007375>, 2007.
- Ciais, P., Sabine, C., Bala, G., Bopp, L., Brovkin, V., Canadell, J., Chhabra, A., DeFries, R., Galloway, J., Heimann, M., Jones, C., Le Quéré, C., Myneni, R., Piao, S., and Thornton, P.: Carbon and Other Biogeochemical Cycles, pp. 465–570, Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, <https://doi.org/10.1017/CBO9781107415324.015>, 2013.
- 985 Ciais, P., Yao, Y., Gasser, T., Baccini, A., Wang, Y., Lauerwald, R., Peng, S., Bastos, A., Li, W., Raymond, P. A., Canadell, J. G., Peters, G. P., Andres, R. J., Chang, J., Yue, C., Dolman, A. J., Haverd, V., Hartmann, J., Laruelle, G., Konings, A. G., King, A. W., Liu, Y., Luysaert, S., Maignan, F., Patra, P. K., Peregon, A., Regnier, P., Pongratz, J., Poulter, B., Shvidenko, A., Valentini, R., Wang, R., Broquet, G., Yin, Y., Zscheischler, J., Guenet, B., Goll, D. S., Ballantyne, A. P., Yang, H., Qiu, C., and Zhu, D.: Empirical estimates of regional carbon budgets imply reduced global soil heterotrophic respiration, *Nat. Sci. Rev.*, 8, nwaal45, <https://doi.org/10.1093/nsr/nwaal45>, 2021.
- 990 Ciais, P., Bastos, A., Chevallier, F., Lauerwald, R., Poulter, B., Canadell, J. G., Hugelius, G., Jackson, R. B., Jain, A., Jones, M., Kondo, M., Luijkx, I. T., Patra, P. K., Peters, W., Pongratz, J., Petrescu, A. M. R., Piao, S., Qiu, C., Von Randow, C., Regnier, P., Saunio, M., Scholes, R., Shvidenko, A., Tian, H., Yang, H., Wang, X., and Zheng, B.: Definitions and methods to estimate regional land carbon fluxes for the second phase of the REgional Carbon Cycle Assessment and Processes Project (RECCAP-2), *Geoscientific Model Development*, 15, 1289–1316, <https://doi.org/10.5194/gmd-15-1289-2022>, 2022.
- 995 Cook-Patton, S. C., Leavitt, S. M., Gibbs, D., Harris, N. L., Lister, K., Anderson-Teixeira, K. J., Briggs, R. D., Chazdon, R. L., Crowther, T. W., Ellis, P. W., Griscom, H. P., Herrmann, V., Holl, K. D., Houghton, R. A., Larrosa, C., Lomax, G., Lucas, R., Madsen, P., Malhi, Y., Paquette, A., Parker, J. D., Paul, K., Routh, D., Roxburgh, S., Saatchi, S., van den Hoogen, J., Walker, W. S., Wheeler, C. E., Wood, S. A., Xu, L., and Griscom, B. W.: Mapping carbon accumulation potential from global natural forest regrowth, *Nature*, 585, 545–550, <https://doi.org/10.1038/s41586-020-2686-x>, 2020.
- 1000 Cressie, N. and Kang, E. L.: Hot enough for you? A spatial exploratory and inferential analysis of North American climate-change projections, *Math. Geosci.*, 48, 107–121, <https://doi.org/10.1007/s11004-015-9607-9>, 2016.
- Crisp, D., Meijer, Y., Munro, R., Bowman, K., Chatterjee, A., Baker, D., Chevallier, F., Nassar, R., Palmer, P. I., Agusti-Panareda, A., Al-Saadi, J., Ariel, Y., Basu, S., Bergamaschi, P., Boesch, H., Bousquet, P., Bovensmann, H., Bréon, F.-M., Brunner, D., Buchwitz, M.,



- 1005 Buisson, F., Burrows, J. P., Butz, A., Ciais, P., Clerbaux, C., Counet, P., Crevoisier, C., Crowell, S., DeCola, P. L., Deniel, C., Dowell, M., Eckman, R., Edwards, D., Ehret, G., Eldering, A., Engelen, R., Fisher, B., Germain, S., Hakkarainen, J., Hilsenrath, E., Holmlund, K., Houweling, S., Hu, H., Jacob, D., Janssens-Maenhout, G., Jones, D. B. A., Jouglet, D., Kataoka, F., Kiel, M., Kulawik, S. S., Kuze, A., Lachance, R. L., Lang, R., Landgraf, J., Liu, J., Liu, Y., Maksyutov, S., Matsunaga, T., McKeever, J., Moore, B., Nakajima, M., Natraj, V., Nelson, R. R., Niwa, Y., Oda, T., O'Dell, C. W., Ott, L., Patra, P., Pawson, S., Payne, V., Pinty, B., Polavarapu, S. M., Retscher, C., Rosenberg, R., Schuh, A., Schwandner, F. M., Shiomi, K., Su, W., Tamminen, J., Taylor, T. E., Veefkind, P., Veihelmann, B., Wofsy, S.,
- 1010 Worden, J., Wunch, D., Yang, D., Zhang, P., and Zehner, C.: A constellation architecture for monitoring carbon dioxide and methane from space, Tech. rep., 32nd Committee on Earth Observation Satellites 2018 Plenary, Brussels, Belgium, October 16-18, 2018, 2018.
- Crisp, D., Dolman, H., Tanhua, T., McKinley, G. A., Hauck, J., Bastos, A., Sitch, S., Eggleston, S., and Aich, V.: How Well Do We Understand the Land-Ocean-Atmosphere Carbon Cycle?, *Rev. Geophys.*, 60, e2021RG000736, <https://doi.org/10.1029/2021RG000736>, 2022.
- 1015 Crowell, S., Baker, D., Schuh, A., Basu, S., Jacobson, A. R., Chevallier, F., Liu, J., Deng, F., Feng, L., McKain, K., Chatterjee, A., Miller, J. B., Stephens, B. B., Eldering, A., Crisp, D., Schimel, D., Nassar, R., O'Dell, C. W., Oda, T., Sweeney, C., Palmer, P. I., and Jones, D. B. A.: The 2015–2016 carbon cycle as seen from OCO-2 and the global in situ network, *Atmos. Chem. Phys.*, 19, 9797–9831, <https://doi.org/10.5194/acp-19-9797-2019>, 2019.
- De Mazière, M., Sha, M. K., Desmet, F., Hermans, C., Scolas, F., Kumps, N., Metzger, J.-M., Duflot, V., and Cammas, J.-P.: TCCON data from Réunion Island (RE), Release GGG2014.R1, <https://doi.org/10.14291/TCCON.GGG2014.REUNION01.R1>, 2017.
- 1020 Deng, F., Jones, D. B. A., Henze, D. K., Bousserez, N., Bowman, K. W., Fisher, J. B., Nassar, R., O'Dell, C., Wunch, D., Wennberg, P. O., Kort, E. A., Wofsy, S. C., Blumenstock, T., Deutscher, N. M., Griffith, D. W. T., Hase, F., Heikkinen, P., Sherlock, V., Strong, K., Sussmann, R., and Warneke, T.: Inferring regional sources and sinks of atmospheric CO<sub>2</sub> from GOSAT XCO<sub>2</sub> data, *Atmos. Chem. Phys.*, 14, 3703–3727, <https://doi.org/10.5194/acp-14-3703-2014>, 2014.
- Deng, F., Jones, D., O'Dell, C. W., Nassar, R., and Parazoo, N. C.: Combining GOSAT XCO<sub>2</sub> observations over land and ocean to improve regional CO<sub>2</sub> flux estimates, *J. Geophys. Res.-Atmos.*, 121, 1896–1913, <https://doi.org/10.1002/2015JD024157>, 2016.
- 1025 Deng, Z., Ciais, P., Tzompa-Sosa, Z. A., Saunio, M., Qiu, C., Tan, C., Sun, T., Ke, P., Cui, Y., Tanaka, K., Lin, X., Thompson, R. L., Tian, H., Yao, Y., Huang, Y., Lauerwald, R., Jain, A. K., Xu, X., Bastos, A., Sitch, S., Palmer, P. I., Lauvaux, T., d'Aspremont, A., Giron, C., Benoit, A., Poulter, B., Chang, J., Petrescu, A. M. R., Davis, S. J., Liu, Z., Grassi, G., Albergel, C., Tubiello, F. N., Perugini, L., Peters, W., and Chevallier, F.: Comparing national greenhouse gas budgets reported in UNFCCC inventories against atmospheric inversions, *Earth*
- 1030 *Syst. Sci. Data*, 14, 1639–1675, <https://doi.org/10.5194/essd-14-1639-2022>, 2022.
- Denning, A. S., Fung, I. Y., and Randall, D.: Latitudinal gradient of atmospheric CO<sub>2</sub> due to seasonal exchange with land biota, *Nature*, 376, 240, <https://doi.org/10.1038/376240a0>, 1995.
- Denning, A. S., Holzer, M., Gurney, K. R., Heimann, M., Law, R. M., Rayner, P. J., Fung, I. Y., Fan, S.-M., Taguchi, S., Friedlingstein, P., Balkanski, Y., Taylor, J., Maiss, M., and Levin, I.: Three-dimensional transport and concentration of SF<sub>2</sub> A model intercomparison study (TransCom 2), *Tellus B*, 51, 266–297, <https://doi.org/10.3402/tellusb.v51i2.16286>, 1999a.
- 1035 Denning, A. S., Takahashi, T., and Friedlingstein, P.: Can a strong atmospheric CO<sub>2</sub> rectifier effect be reconciled with a “reasonable” carbon budget?, *Tellus B*, 51, 249–253, <https://doi.org/10.3402/tellusb.v51i2.16277>, 1999b.
- Denvil-Sommer, A., Gehlen, M., Vrac, M., and Mejia, C.: LSCE-FFNN-v1: a two-step neural network model for the reconstruction of surface ocean pCO<sub>2</sub> over the global ocean, *Geosci. Model Dev.*, 12, 2091–2105, <https://doi.org/10.5194/gmd-12-2091-2019>, 2019.



- 1040 Dubey, M., Henderson, B., Green, D., Butterfield, Z., Keppel-Aleks, G., Allen, N., Blavier, J.-F., Roehl, C., Wunch, D., and Lindenmaier, R.: TCCON data from Manaus (BR), Release GGG2014R0, TCCON data archive, hosted by CaltechDATA, <https://doi.org/10.14291/tcon.ggg2014.manau01.R0/1149274>, 2014.
- Enting, I. and Mansbridge, J.: Latitudinal distribution of sources and sinks of CO<sub>2</sub>: Results of an inversion study, *Tellus B*, 43, 156–170, <https://doi.org/10.3402/tellusb.v43i2.15261>, 1991.
- 1045 Enting, I., Trudinger, C., and Francey, R.: A synthesis inversion of the concentration and  $\delta^{13}\text{C}$  of atmospheric CO<sub>2</sub>, *Tellus B*, 47, 35–52, <https://doi.org/10.1034/j.1600-0889.47.issue1.5.x>, 1995.
- EPA: Inventory of U.S. Greenhouse Gas Emissions and Sinks: 1990-2020. U.S. Environmental Protection Agency, EPA 430-R-22-003, <https://www.epa.gov/ghgemissions/draft-inventory-us-greenhouse-gas-emissions-and-sinks-1990-2020>, 2022.
- Fay, A. R. and McKinley, G. A.: Observed Regional Fluxes to Constrain Modeled Estimates of the Ocean Carbon Sink, *Geophys. Res. Lett.*, 48, e2021GL095325, <https://doi.org/10.1029/2021GL095325>, 2021.
- 1050 Fay, A. R., Gregor, L., Landschützer, P., McKinley, G. A., Gruber, N., Gehlen, M., Iida, Y., Laruelle, G. G., Rödenbeck, C., Roobaert, A., and Zeng, J.: SeaFlux: harmonization of air–sea CO<sub>2</sub> fluxes from surface *p*CO<sub>2</sub> data products using a standardized approach, *Earth Syst. Sci. Data*, 13, 4693–4710, <https://doi.org/10.5194/essd-13-4693-2021>, 2021.
- Feist, D. G., Arnold, S. G., John, N., and Geibel, M. C.: TCCON data from Ascension Island (SH), Release GGG2014R0, TCCON data archive, hosted by CaltechDATA, <https://doi.org/10.14291/tcon.ggg2014.ascension01.R0/1149285>, 2014.
- 1055 Feng, L., Palmer, P. I., Parker, R. J., Deutscher, N. M., Feist, D. G., Kivi, R., Morino, I., and Sussmann, R.: Estimates of European uptake of CO<sub>2</sub> inferred from GOSAT X<sub>CO<sub>2</sub></sub> retrievals: sensitivity to measurement bias inside and outside Europe, *Atmos. Chem. Phys.*, 16, 1289–1302, <https://doi.org/10.5194/acp-16-1289-2016>, 2016.
- Feng, S., Lauvaux, T., Davis, K. J., Keller, K., Zhou, Y., Williams, C., Schuh, A. E., Liu, J., and Baker, I.: Seasonal characteristics of model uncertainties from biogenic fluxes, transport, and large-scale boundary inflow in atmospheric CO<sub>2</sub> simulations over North America, *J. Geophys. Res.-Atmos.*, 124, 14 325–14 346, <https://doi.org/10.1029/2019JD031165>, 2019.
- 1060 Frank, D., Reichstein, M., Bahn, M., Thonicke, K., Frank, D., Mahecha, M. D., Smith, P., Van der Velde, M., Vicca, S., Babst, F., Beer, C., Buchmann, N., Canadell, J. G., Ciais, P., Cramer, W., Ibrom, A., Miglietta, F., Poulter, B., Rammig, A., Seneviratne, S. I., Walz, A., Wattenbach, M., Zavala, M. A., and Zscheischler, J.: Effects of climate extremes on the terrestrial carbon cycle: concepts, processes and potential future impacts, *Global Change Biology*, 21, 2861–2880, <https://doi.org/10.1111/gcb.12916>, 2015.
- 1065 Frey, M., Sha, M. K., Hase, F., Kiel, M., Blumenstock, T., Harig, R., Surawicz, G., Deutscher, N. M., Shiomi, K., Franklin, J. E., Bösch, H., Chen, J., Grutter, M., Ohshima, H., Sun, Y., Butz, A., Mengistu Tsidu, G., Ene, D., Wunch, D., Cao, Z., Garcia, O., Ramonet, M., Vogel, F., and Orphal, J.: Building the Collaborative Carbon Column Observing Network (COCCON): long-term stability and ensemble performance of the EM27/SUN Fourier transform spectrometer, *Atmos. Meas. Tech.*, 12, 1513–1530, <https://doi.org/10.5194/amt-12-1513-2019>, 2019.
- 1070 Friedlingstein, P., Jones, M. W., O’Sullivan, M., Andrew, R. M., Bakker, D. C. E., Hauck, J., Le Quééré, C., Peters, G. P., Peters, W., Pongratz, J., Sitch, S., Canadell, J. G., Ciais, P., Jackson, R. B., Alin, S. R., Anthoni, P., Bates, N. R., Becker, M., Bellouin, N., Bopp, L., Chau, T. T. T., Chevallier, F., Chini, L. P., Cronin, M., Currie, K. I., Decharme, B., Djeutchouang, L. M., Dou, X., Evans, W., Feely, R. A., Feng, L., Gasser, T., Gilfillan, D., Gkritzalis, T., Grassi, G., Gregor, L., Gruber, N., Gürses, O., Harris, I., Houghton, R. A., Hurtt, G. C., Iida, Y., Ilyina, T., Luijkx, I. T., Jain, A., Jones, S. D., Kato, E., Kennedy, D., Klein Goldewijk, K., Knauer, J., Korsbakken, J. I., Körtzinger, A., Landschützer, P., Lauvset, S. K., Lefèvre, N., Lienert, S., Liu, J., Marland, G., McGuire, P. C., Melton, J. R., Munro, D. R., Nabel, J. E. M. S., Nakaoka, S.-I., Niwa, Y., Ono, T., Pierrot, D., Poulter, B., Rehder, G., Resplandy, L., Robertson, E., Rödenbeck, C., Rosan,



- 1080 T. M., Schwinger, J., Schwingshackl, C., Séférian, R., Sutton, A. J., Sweeney, C., Tanhua, T., Tans, P. P., Tian, H., Tilbrook, B., Tubiello, F., van der Werf, G. R., Vuichard, N., Wada, C., Wanninkhof, R., Watson, A. J., Willis, D., Wiltshire, A. J., Yuan, W., Yue, C., Yue, X., Zaehle, S., and Zeng, J.: Global Carbon Budget 2021, *Earth Syst. Sci. Data*, 14, 1917–2005, <https://doi.org/10.5194/essd-14-1917-2022>, 2022.
- Gatti, L. V., Basso, L. S., Miller, J. B., Gloor, M., Gatti Domingues, L., Cassol, H. L., Tejada, G., Aragão, L. E., Nobre, C., Peters, W., Marani, L., Arai, E., Sanches, A. H., Corrêa, S. M., Anderson, L., Rando, C. V., Correia, C. S. C., Crispim, S. P., and Neves, R. A. L.: Amazonia as a carbon source linked to deforestation and climate change, *Nature*, 595, 388–393, <https://doi.org/10.1038/s41586-021-03629-6>, 2021.
- 1085 Gaubert, B., Stephens, B. B., Basu, S., Chevallier, F., Deng, F., Kort, E. A., Patra, P. K., Peters, W., Rödenbeck, C., Saeki, T., Schimel, D., Van der Laan-Luijkx, I., Wofsy, S., and Yin, Y.: Global atmospheric CO<sub>2</sub> inverse models converging on neutral tropical land exchange, but disagreeing on fossil fuel and atmospheric growth rate, *Biogeosciences*, 16, 117–134, <https://doi.org/10.5194/bg-16-117-2019>, 2019.
- Girard, C., Plante, A., Desgagné, M., McTaggart-Cowan, R., Côté, J., Charron, M., Gravel, S., Lee, V., Patoine, A., Qaddouri, A., Roch, M., Spacek, L., Tanguay, M., Vaillancourt, P. A., and Zadra, A.: Staggered vertical discretization of the Canadian Environmental Multiscale (GEM) model using a coordinate of the log-hydrostatic-pressure type, *Mon. Weather Rev.*, 142, 1183–1196, <https://doi.org/10.1175/MWR-D-13-00255.1>, 2014.
- 1090 Gloor, M., Bakwin, P., Hurst, D., Lock, L., Draxler, R., and Tans, P.: What is the concentration footprint of a tall tower?, *J. Geophys. Res.-Atmos.*, 106, 17 831–17 840, <https://doi.org/10.1029/2001JD900021>, 2001.
- Goo, T.-Y., Oh, Y.-S., and Velazco, V. A.: TCCON data from Anmeyondo (KR), Release GGG2014R0, TCCON data archive, hosted by CaltechDATA, <https://doi.org/10.14291/tcon.ggg2014.anmeyondo01.R0/1149284>, 2014.
- 1095 Grassi, G., House, J., Kurz, W. A., Cescatti, A., Houghton, R. A., Peters, G. P., Sanz, M. J., Viñas, R. A., Alkama, R., Arneeth, A., Bondeau, A., Dentener, F., Fader, M., Federici, S., Friedlingstein, P., Jain, A. K., Kato, E., Koven, C. D., Lee, D., Nabel, J. E. M. S., Nassikas, A. A., Perugini, L., Rossi, S., Sitch, S., Viovy, N., Wiltshire, A., and Zaehle, S.: Reconciling global-model estimates and country reporting of anthropogenic forest CO<sub>2</sub> sinks, *Nat. Clim. Change*, 8, 914–920, <https://doi.org/10.1038/s41558-018-0283-x>, 2018.
- 1100 Gregor, L. and Fay, A.: SeaFlux: Air-sea CO<sub>2</sub> fluxes for surface pCO<sub>2</sub> data products using a standardised approach, <https://doi.org/10.5281/zenodo.5078404>, version 2021.02, 2021.
- Gregor, L., Lebehot, A. D., Kok, S., and Scheel Monteiro, P. M.: A comparative assessment of the uncertainties of global surface ocean CO<sub>2</sub> estimates using a machine-learning ensemble (CSIR-ML6 version 2019a) – have we hit the wall?, *Geoscientific Model Development*, 12, 5113–5136, <https://doi.org/10.5194/gmd-12-5113-2019>, 2019.
- 1105 Griffith, D. W., Deutscher, N. M., Velazco, V. A., Wennberg, P. O., Yavin, Y., Aleks, G. K., Washenfelder, R. a., Toon, G. C., Blavier, J.-F., Murphy, C., Jones, N., Kettlewell, G., Connor, B. J., Macatangay, R., Roehl, C., Ryzcek, M., Glowacki, J., Culf, T., and Bryant, G.: TCCON data from Darwin (AU), Release GGG2014R0, TCCON data archive, hosted by CaltechDATA, <https://doi.org/10.14291/tcon.ggg2014.darwin01.R0/1149290>, 2014a.
- Griffith, D. W., Velazco, V. A., Deutscher, N. M., Murphy, C., Jones, N., Wilson, S., Macatangay, R., Kettlewell, G., Buchholz, R. R., and Riggenschach, M.: TCCON data from Wollongong (AU), Release GGG2014R0, TCCON data archive, hosted by CaltechDATA, <https://doi.org/10.14291/tcon.ggg2014.wollongong01.R0/1149291>, 2014b.
- 1110 Guan, D., Liu, Z., Geng, Y., Lindner, S., and Hubacek, K.: The gigatonne gap in China’s carbon dioxide inventories, *Nat. Clim. Change*, 2, 672–675, <https://doi.org/10.1038/nclimate1560>, 2012.
- Gurney, K. R., Law, R. M., Denning, A. S., Rayner, P. J., Baker, D., Bousquet, P., Bruhwiler, L., Chen, Y.-H., Ciais, P., Fan, S., Fung, I. Y., Gloor, M., Heimann, M., Higuchi, K., John, J., Maki, T., Maksyutov, S., Masarie, K., Peylin, P., Prather, M., Pak, B. C., Randerson, J.,





- Sarmiento, J., Taguchi, S., Takahashi, T., and Yuen, C.-W.: Towards robust regional estimates of CO<sub>2</sub> sources and sinks using atmospheric transport models, *Nature*, 415, 626–630, <https://doi.org/10.1038/415626a>, 2002.
- Hall, B. D., Crotwell, A. M., Kitzis, D. R., Mefford, T., Miller, B. R., Schibig, M. F., and Tans, P. P.: Revision of the World Meteorological Organization Global Atmosphere Watch (WMO/GAW) CO<sub>2</sub> calibration scale, *Atmos. Meas. Tech.*, 14, 3015–3032, <https://doi.org/10.5194/amt-14-3015-2021>, 2021.
- 1120 Harris, N. L., Brown, S., Hagen, S. C., Saatchi, S. S., Petrova, S., Salas, W., Hansen, M. C., Potapov, P. V., and Lotsch, A.: Baseline map of carbon emissions from deforestation in tropical regions, *Science*, 336, 1573–1576, <https://doi.org/10.1126/science.1217962>, 2012.
- Hase, F., Blumenstock, T., Dohe, S., Gross, J., and Kiel, M.: TCCON data from Karlsruhe (DE), Release GGG2014R1, TCCON data archive, hosted by CaltechDATA, <https://doi.org/10.14291/tcon.ggg2014.karlsruhe01.R1/1182416>, 2014.
- 1125 Hu, L., Andrews, A. E., Thoning, K. W., Sweeney, C., Miller, J. B., Michalak, A. M., Dlugokencky, E., Tans, P. P., Shiga, Y. P., Mountain, M., Nehrkorn, T., Montzka, S. A., McKain, K., Kofler, J., Trudeau, M., Michel, S. E., Biraud, S. C., Fischer, M. L., Worthy, D. E. J., Vaughn, B. H., White, J. W. C., Yadav, V., Basu, S., and van der Velde, I. R.: Enhanced North American carbon uptake associated with El Niño, *Science advances*, 5, eaaw0076, <https://doi.org/10.1126/sciadv.aaw0076>, 2019.
- Hu, L., Montzka, S. A., Kaushik, A., Andrews, A. E., Sweeney, C., Miller, J., Baker, I. T., Denning, S., Campbell, E., Shiga, Y. P., Tans, P., Siso, M. C., Crotwell, M., McKain, K., Thoning, K., Hall, B., Vimont, I., Elkins, J. W., Whelan, M. E., and Suntharalingam, P.: COS-derived GPP relationships with temperature and light help explain high-latitude atmospheric CO<sub>2</sub> seasonal cycle amplification, *Proceedings of the National Academy of Sciences*, 118, e2103423 118, <https://doi.org/10.1073/pnas.2103423118>, 2021.
- 1130 Iida, Y., Takatani, Y., Kojima, A., and Ishii, M.: Global trends of ocean CO<sub>2</sub> sink and ocean acidification: an observation-based reconstruction of surface ocean inorganic carbon variables, *J. Oceanogr.*, 77, 323–358, <https://doi.org/10.1007/s10872-020-00571-5>, 2021.
- 1135 IPCC: 2006 IPCC Guidelines for National Greenhouse Gas Inventories, Prepared by the National Greenhouse Gas Inventories Programme, IGES, Japan, 2006.
- IPCC: 2019 Refinement to the 2006 IPCC Guidelines for National Greenhouse Gas Inventories, IPCC, Switzerland, 2019.
- Iraci, L. T., Podolske, J., Hillyard, P. W., Roehl, C., Wennberg, P. O., Blavier, J.-F., Allen, N., Wunch, D., Osterman, G. B., and Albertson, R.: TCCON data from Edwards (US), Release GGG2014R1, TCCON data archive, hosted by CaltechDATA, <https://doi.org/10.14291/tcon.ggg2014.edwards01.R1/1255068>, 2016.
- 1140 Jacobson, A. R., Schuldt, K. N., Miller, J. B., Oda, T., Tans, P., Arlyn Andrews, Mund, J., Ott, L., Collatz, G. J., Aalto, T., Afshar, S., Aikin, K., Aoki, S., Apadula, F., Baier, B., Bergamaschi, P., Beyersdorf, A., Biraud, S. C., Bollenbacher, A., Bowling, D., Brailsford, G., Abshire, J. B., Chen, G., Huilin Chen, Lukasz Chmura, Sites Climadat, Colomb, A., Conil, S., Cox, A., Cristofanelli, P., Cuevas, E., Curcoll, R., Sloop, C. D., Davis, K., Wekker, S. D., Delmotte, M., DiGangi, J. P., Dlugokencky, E., Ehleringer, J., Elkins, J. W.,
- 1145 Emmenegger, L., Fischer, M. L., Forster, G., Frumau, A., Galkowski, M., Gatti, L. V., Gloor, E., Griffis, T., Hammer, S., Haszpra, L., Hatakka, J., Heliasz, M., Hensen, A., Hermanssen, O., Hintsa, E., Holst, J., Jaffe, D., Karion, A., Kawa, S. R., Keeling, R., Keronen, P., Kolari, P., Kominkova, K., Kort, E., Krummel, P., Kubistin, D., Labuschagne, C., Langenfelds, R., Laurent, O., Laurila, T., Lauvaux, T., Law, B., Lee, J., Lehner, I., Leuenberger, M., Levin, I., Levula, J., Lin, J., Lindauer, M., Loh, Z., Lopez, M., Luijkx, I. T., Myhre, C. L., Machida, T., Mammarella, I., Manca, G., Manning, A., Manning, A., Marek, M. V., Marklund, P., Martin, M. Y., Matsueda, H.,
- 1150 McKain, K., Meijer, H., Meinhardt, F., Miles, N., Miller, C. E., Mölder, M., Montzka, S., Moore, F., Josep-Anton Morgui, Morimoto, S., Munger, B., Jaroslaw Necki, Newman, S., Nichol, S., Niwa, Y., O’Doherty, S., Mikael Ottosson-Löfvenius, Paplawsky, B., Peischl, J., Peltola, O., Jean-Marc Pichon, Piper, S., Plass-Dömler, C., Ramonet, M., Reyes-Sanchez, E., Richardson, S., Riris, H., Ryerson, T., Saito, K., Sargent, M., Sasakawa, M., Sawa, Y., Say, D., Scheeren, B., Schmidt, M., Schmidt, A., Schumacher, M., Shepson, P., Shook,



- M., Stanley, K., Steinbacher, M., Stephens, B., Sweeney, C., Thoning, K., Torn, M., Turnbull, J., Tørseth, K., Bulk, P. V. D., Dincer, D. V., Vermeulen, A., Viner, B., Vitkova, G., Walker, S., Weyrauch, D., Wofsy, S., Worthy, D., Dickon Young, and Mirosław Zimnoch: CarbonTracker CT2019B, <https://doi.org/10.25925/20201008>, 2020.
- Janssens-Maenhout, G., Pinty, B., Dowell, M., Zunker, H., Andersson, E., Balsamo, G., Bézy, J.-L., Brunhes, T., Bösch, H., Bojkov, B., Brunner, D., Buchwitz, M., Crisp, D., Ciais, P., Counet, P., Dee, D., Denier van der Gon, H., Dolman, H., Drinkwater, M. R., Dubovik, O., Engelen, R., Fehr, T., Fernandez, V., Heimann, M., Holmlund, K., Houweling, S., Husband, R., Juvyns, O., Kentarchos, A., Landgraf, J., Lang, R., Löscher, A., Marshall, J., Meijer, Y., Nakajima, M., Palmer, P. I., Peylin, P., Rayner, P., Scholze, M., Sierk, B., Tamminen, J., and Veefkind, P.: Toward an Operational Anthropogenic CO<sub>2</sub> Emissions Monitoring and Verification Support Capacity, *B. A. Meteorol. Soc.*, 101, E1439–E1451, <https://doi.org/10.1175/BAMS-D-19-0017.1>, 2020.
- Karion, A., Sweeney, C., Tans, P., and Newberger, T.: AirCore: An innovative atmospheric sampling system, *J. Atmos Ocean Tech.*, 27, 1839–1853, <https://doi.org/10.1175/2010JTECHA1448.1>, 2010.
- Kaushik, A., Graham, J., Dorheim, K. R., Kramer, R., Wang, J., and Byrne, B.: The future of the carbon cycle in a changing climate, *Eos*, 101, <https://doi.org/https://doi.org/10.1029/2020EO140276>, 2020.
- Kawakami, S., Ohyama, H., Arai, K., Okumura, H., Taura, C., Fukamachi, T., and Sakashita, M.: TCCON data from Saga (JP), Release GGG2014R0, TCCON data archive, hosted by CaltechDATA, <https://doi.org/10.14291/tcon.ggg2014.saga01.R0/1149283>, 2014.
- Keeling, R. F. and Graven, H. D.: Insights from Time Series of Atmospheric Carbon Dioxide and Related Tracers, *Annual Review of Environment and Resources*, 46, 85–110, <https://doi.org/10.1146/annurev-environ-012220-125406>, 2021.
- Keeling, R. F., Graven, H. D., Welp, L. R., Resplandy, L., Bi, J., Piper, S. C., Sun, Y., Bollenbacher, A., and Meijer, H. A.: Atmospheric evidence for a global secular increase in carbon isotopic discrimination of land photosynthesis, *Proceedings of the National Academy of Sciences*, 114, 10361–10366, 2017.
- Kiel, M., O'Dell, C. W., Fisher, B., Eldering, A., Nassar, R., MacDonald, C. G., and Wennberg, P. O.: How bias correction goes wrong: measurement of X<sub>CO<sub>2</sub></sub> affected by erroneous surface pressure estimates, *Atmos. Meas. Tech.*, 12, 2241–2259, <https://doi.org/10.5194/amt-12-2241-2019>, 2019.
- Kivi, R., Heikkinen, P., and Kyrö, E.: TCCON data from Sodankyla (FI), Release GGG2014R0, TCCON data archive, hosted by CaltechDATA, <https://doi.org/10.14291/tcon.ggg2014.sodankyla01.R0/1149280>, 2014.
- Kondo, M., Ichii, K., Patra, P. K., Poulter, B., Calle, L., Koven, C., Pugh, T. A., Kato, E., Harper, A., Zaehle, S., et al.: Plant regrowth as a driver of recent enhancement of terrestrial CO<sub>2</sub> uptake, *Geophys. Res. Lett.*, 45, 4820–4830, 2018.
- Kondo, M., Patra, P. K., Sitch, S., Friedlingstein, P., Poulter, B., Chevallier, F., Ciais, P., Canadell, J. G., Bastos, A., Lauerwald, R., et al.: State of the science in reconciling top-down and bottom-up approaches for terrestrial CO<sub>2</sub> budget, *Global change biology*, 26, 1068–1084, <https://doi.org/10.1111/gcb.14917>, 2020.
- Krol, M., de Bruine, M., Killaars, L., Ouwersloot, H., Pozzer, A., Yin, Y., Chevallier, F., Bousquet, P., Patra, P., Belikov, D., Maksyutov, S., Dhomse, S., Feng, W., and Chipperfield, M. P.: Age of air as a diagnostic for transport timescales in global models, *Geosci. Model Dev.*, 11, 3109–3130, <https://doi.org/10.5194/gmd-11-3109-2018>, 2018.
- Landschützer, P., Gruber, N., Bakker, D. C., and Schuster, U.: Recent variability of the global ocean carbon sink, *Global Biogeochem. Cy.*, 28, 927–949, <https://doi.org/10.1002/2014GB004853>, 2014.
- Landschützer, P., Gruber, N., and Bakker, D. C. E.: An observation-based global monthly gridded sea surface pCO<sub>2</sub> product from 1982 onward and its monthly climatology (NCEI Accession 0160558). Version 5.5., <https://doi.org/10.7289/V5Z899N6>, 2020.



- Law, R. and Simmonds, I.: The sensitivity of deduced CO<sub>2</sub> sources and sinks to variations in transport and imposed surface concentrations, *Tellus B*, 48, 613–625, <https://doi.org/10.1034/j.1600-0889.1996.t01-4-00001.x>, 1996.
- Law, R., Rayner, P., Denning, A., Erickson, D., Fung, I., Heimann, M., Piper, S., Ramonet, M., Taguchi, S., Taylor, J., et al.: Variations in modeled atmospheric transport of carbon dioxide and the consequences for CO<sub>2</sub> inversions, *Global Biogeochem. Cy.*, 10, 783–796, <https://doi.org/10.1029/96GB01892>, 1996.
- 1195
- Lawson, C. L. and Hanson, R.: Linear least squares with linear inequality constraints, *Solving least squares problems*, pp. 158–173, 1974.
- Liu, C., Wang, W., and Sun, Y.: TCCON data from Hefei(PRC), Release GGG2014.R0, TCCON data archive, hosted by CaltechDATA, <https://doi.org/10.14291/tcon.ggg2014.hefei01.R0>, 2018.
- Liu, J., Fung, I., Kalnay, E., and Kang, J.-S.: CO<sub>2</sub> transport uncertainties from the uncertainties in meteorological fields, *Geophys. Res. Lett.*, 38, <https://doi.org/10.1029/2011GL047213>, 2011.
- 1200
- Liu, J., Bowman, K. W., and Henze, D. K.: Source-receptor relationships of column-average CO<sub>2</sub> and implications for the impact of observations on flux inversions, *J. Geophys. Res.-Atmos.*, 120, 5214–5236, <https://doi.org/10.1002/2014JD022914>, 2015.
- Liu, J., Bowman, K. W., Schimel, D. S., Parazoo, N. C., Jiang, Z., Lee, M., Bloom, A. A., Wunch, D., Frankenberg, C., Sun, Y., O'Dell, C. W., Gurney, K. R., Menemenlis, D., Gierach, M., Crisp, D., and Eldering, A.: Contrasting carbon cycle responses of the tropical continents to the 2015–2016 El Niño, *Science*, 358, <https://doi.org/10.1126/science.aam5690>, 2017.
- 1205
- Liu, J., Baskaran, L., Bowman, K., Schimel, D., Bloom, A. A., Parazoo, N. C., Oda, T., Carroll, D., Menemenlis, D., Joiner, J., Commane, R., Daube, B., Gatti, L. V., McKain, K., Miller, J., Stephens, B. B., Sweeney, C., and Wofsy, S.: Carbon Monitoring System Flux Net Biosphere Exchange 2020 (CMS-Flux NBE 2020), *Earth Syst. Sci. Data*, 13, 299–330, <https://doi.org/10.5194/essd-13-299-2021>, 2021a.
- Liu, M., Tian, H., Yang, Q., Yang, J., Song, X., Lohrenz, S. E., and Cai, W.-J.: Long-term trends in evapotranspiration and runoff over the drainage basins of the Gulf of Mexico during 1901–2008, *Water Resour. Res.*, 49, 1988–2012, <https://doi.org/10.1002/wrcr.20180>, 2013.
- 1210
- Liu, Z., Zeng, N., Liu, Y., Kalnay, E., Asrar, G., Wu, B., Cai, Q., Liu, D., and Han, P.: Improving the joint estimation of CO<sub>2</sub> and surface carbon fluxes using a Constrained Ensemble Kalman Filter in COLA (v1.0), *Geosci. Model Dev. Discussions*, 2021, 1–25, <https://doi.org/10.5194/gmd-2021-375>, 2021b.
- Maksyutov, S., Oda, T., Saito, M., Janardanan, R., Belikov, D., Kaiser, J. W., Zhuravlev, R., Ganshin, A., Valsala, V. K., Andrews, A., Chmura, L., Dlugokencky, E., Haszpra, L., Langenfelds, R. L., Machida, T., Nakazawa, T., Ramonet, M., Sweeney, C., and Worthy, D.: Technical note: A high-resolution inverse modelling technique for estimating surface CO<sub>2</sub> fluxes based on the NIES-TM-FLEXPART coupled transport model and its adjoint, *Atmos. Chem. Phys.*, 21, 1245–1266, <https://doi.org/10.5194/acp-21-1245-2021>, 2021.
- 1215
- Masarie, K., Peters, W., Jacobson, A., and Tans, P.: ObsPack: a framework for the preparation, delivery, and attribution of atmospheric greenhouse gas measurements, *Earth Syst. Sci. Data*, 6, 375–384, <https://doi.org/10.5194/essd-6-375-2014>, 2014.
- 1220
- Mason Earles, J., Yeh, S., and Skog, K. E.: Timing of carbon emissions from global forest clearance, *Nat. Clim. Change*, 2, 682–685, <https://doi.org/10.1038/nclimate1535>, 2012.
- Mayorga, E., Seitzinger, S. P., Harrison, J. A., Dumont, E., Beusen, A. H., Bouwman, A., Fekete, B. M., Kroeze, C., and Van Drecht, G.: Global nutrient export from WaterSheds 2 (NEWS 2): model development and implementation, *Environ. Modell. Softw.*, 25, 837–853, <https://doi.org/10.1016/j.envsoft.2010.01.007>, 2010.
- 1225
- McKinley, G. A., Fay, A. R., Lovenduski, N. S., and Pilcher, D. J.: Natural variability and anthropogenic trends in the ocean carbon sink, *Annu. Rev. Mar. Sci.*, 9, 125–150, <https://doi.org/10.1146/annurev-marine-010816-060529>, 2017.



- McNorton, J. R., Bousserez, N., Agustí-Panareda, A., Balsamo, G., Choulga, M., Dawson, A., Engelen, R., Kipling, Z., and Lang, S.: Representing model uncertainty for global atmospheric CO<sub>2</sub> flux inversions using ECMWF-IFS-46R1, *Geosci. Model Dev.*, 13, 2297–2313, <https://doi.org/10.5194/gmd-13-2297-2020>, 2020.
- 1230 Mearns, L. and et al.: The North American Regional Climate Change Assessment Program dataset, <https://doi.org/10.5065/D6RN35ST>, 2007.
- Messerschmidt, J., Geibel, M. C., Blumenstock, T., Chen, H., Deutscher, N. M., Engel, A., Feist, D. G., Gerbig, C., Gisi, M., Hase, F., Katrynski, K., Kolle, O., Lavrič, J. V., Notholt, J., Palm, M., Ramonet, M., Rettinger, M., Schmidt, M., Sussmann, R., Toon, G. C., Truong, F., Warneke, T., Wennberg, P. O., Wunch, D., and Xueref-Remy, I.: Calibration of TCCON column-averaged CO<sub>2</sub>: the first  
1235 aircraft campaign over European TCCON sites, *Atmos. Chem. Phys.*, 11, 10 765–10 777, <https://doi.org/10.5194/acp-11-10765-2011>, 2011.
- Meybeck, M., Dürr, H. H., and Vörösmarty, C. J.: Global coastal segmentation and its river catchment contributors: A new look at land-ocean linkage, *Global Biogeochem. Cy.*, 20, <https://doi.org/10.1029/2005GB002540>, 2006.
- Miller, C., Crisp, D., DeCola, P., Olsen, S., Randerson, J. T., Michalak, A. M., Alkhaled, A., Rayner, P., Jacob, D. J., Suntharalingam, P.,  
1240 et al.: Precision requirements for space-based X<sub>CO<sub>2</sub></sub> data, *J. Geophys. Res.-Atmos.*, 112, <https://doi.org/10.1029/2006JD007659>, 2007.
- Miller, J. B., Martins, G. A., de Souza, R. A., and Schuldt, K. N.: Manaus Aircraft profile data for the period 2017-2020; obspack\_multi-species\_1\_manauas\_profiles\_v1.0\_2021-05-19, <https://doi.org/10.25925/20210519>, 2021.
- Miller, S. M., Saibaba, A. K., Trudeau, M. E., Mountain, M. E., and Andrews, A. E.: Geostatistical inverse modeling with very large datasets: an example from the Orbiting Carbon Observatory 2 (OCO-2) satellite, *Geosci. Model Dev.*, 13, 1771–1785, <https://doi.org/10.5194/gmd-13-1771-2020>, 2020.  
1245
- Morino, I., Yokozeki, N., Matzuzaki, T., and Horikawa, M.: TCCON data from Rikubetsu (JP), Release GGG2014R1, TCCON data archive, hosted by CaltechDATA, <https://doi.org/10.14291/tcon.ggg2014.rikubetsu01.R1/1242265>, 2014.
- Morino, I., Matsuzaki, T., and Horikawa, M.: TCCON data from Tsukuba (JP), 125HR, Release GGG2014.R2, <https://doi.org/10.14291/TCCON.GGG2014.TSUKUBA02.R2>, 2018a.
- 1250 Morino, I., Velazco, V. A., Akihiro, H., Osamu, U., and Griffith, D. W. T.: TCCON data from Burgos, Ilocos Norte (PH), Release GGG2014.R0, TCCON data archive, hosted by CaltechDATA, <https://doi.org/10.14291/tcon.ggg2014.burgos01.R0>, 2018b.
- Nara, H., Tanimoto, H., Tohjima, Y., Mukai, H., Nojiri, Y., and Machida, T.: Emission factors of CO<sub>2</sub>, CO and CH<sub>4</sub> from Sumatran peatland fires in 2013 based on shipboard measurements, *Tellus B*, 69, 1399 047, <https://doi.org/10.1080/16000889.2017.1399047>, 2017.
- Nassar, R., Jones, D. B. A., Suntharalingam, P., Chen, J. M., Andres, R. J., Wecht, K. J., Yantosca, R. M., Kulawik, S. S., Bowman, K. W.,  
1255 Worden, J. R., Machida, T., and Matsueda, H.: Modeling global atmospheric CO<sub>2</sub> with improved emission inventories and CO<sub>2</sub> production from the oxidation of other carbon species, *Geosci. Model Dev.*, 3, 689, <https://doi.org/10.5194/gmd-3-689-2010>, 2010.
- Nassar, R., Napier-Linton, L., Gurney, K. R., Andres, R. J., Oda, T., Vogel, F. R., and Deng, F.: Improving the temporal and spatial distribution of CO<sub>2</sub> emissions from global fossil fuel emission data sets, *J. Geophys. Res.-Atmos.*, 118, 917–933, <https://doi.org/10.1029/2012JD018196>, 2013.
- 1260 Niwa, Y., Fujii, Y., Sawa, Y., Iida, Y., Ito, A., Satoh, M., Imasu, R., Tsuboi, K., Matsueda, H., and Saigusa, N.: A 4D-Var inversion system based on the icosahedral grid model (NICAM-TM 4D-Var v1.0) – Part 2: Optimization scheme and identical twin experiment of atmospheric CO<sub>2</sub> inversion, *Geosci. Model Dev.*, 10, 2201–2219, <https://doi.org/10.5194/gmd-10-2201-2017>, 2017.
- Notholt, J., Petri, C., Warneke, T., Deutscher, N. M., Palm, M., Buschmann, M., Weinzierl, C., Macatangay, R. C., and Grupe, P.: TCCON data from Bremen (DE), Release GGG2014.R1, <https://doi.org/10.14291/TCCON.GGG2014.BREMEN01.R1>, 2019a.



- 1265 Notholt, J., Warneke, T., Petri, C., Deutscher, N. M., Weinzierl, C., Palm, M., and Buschmann, M.: TCCON data from Ny Ålesund, Spitsbergen (NO), Release GGG2014.R1, <https://doi.org/10.14291/TCCON.GGG2014.NYALESUND01.R1>, 2019b.
- Oda, T. and Maksyutov, S.: A very high-resolution (1 km × 1 km) global fossil fuel CO<sub>2</sub> emission inventory derived using a point source database and satellite observations of nighttime lights, *Atmos. Chem. Phys.*, 11, 543–556, <https://doi.org/10.5194/acp-11-543-2011>, 2011.
- Oda, T., Ott, L., Topylko, P., Halushchak, M., Bun, R., Lesiv, M., Danylo, O., and Horabik-Pyzel, J.: Uncertainty associated with fossil fuel  
1270 carbon dioxide (CO<sub>2</sub>) gridded emission datasets, 2015.
- Oda, T., Maksyutov, S., and Andres, R. J.: The Open-source Data Inventory for Anthropogenic CO<sub>2</sub>, version 2016 (ODIAC2016): a global monthly fossil fuel CO<sub>2</sub> gridded emissions data product for tracer transport simulations and surface flux inversions, *Earth Syst. Sci. Data*, 10, 87–107, <https://doi.org/10.5194/essd-10-87-2018>, 2018.
- Oda, T., Bun, R., Kinakh, V., Topylko, P., Halushchak, M., Marland, G., Lauvaux, T., Jonas, M., Maksyutov, S., Nahorski, Z., et al.: Errors and  
1275 uncertainties in a gridded carbon dioxide emissions inventory, *Mitig. Adapt. Strat. Gl.*, 24, 1007–1050, <https://doi.org/10.1007/s11027-019-09877-2>, 2019.
- O'Dell, C. W., Eldering, A., Wennberg, P. O., Crisp, D., Gunson, M. R., Fisher, B., Frankenberg, C., Kiel, M., Lindqvist, H., Mandrake, L., Merrelli, A., Natraj, V., Nelson, R. R., Osterman, G. B., Payne, V. H., Taylor, T. E., Wunch, D., Drouin, B. J., Oyafuso, F., Chang, A., McDuffie, J., Smyth, M., Baker, D. F., Basu, S., Chevallier, F., Crowell, S. M. R., Feng, L., Palmer, P. I., Dubey, M., García, O. E., Griffith,  
1280 D. W. T., Hase, F., Iraci, L. T., Kivi, R., Morino, I., Notholt, J., Ohyama, H., Petri, C., Roehl, C. M., Sha, M. K., Strong, K., Sussmann, R., Te, Y., Uchino, O., and Velasco, V. A.: Improved retrievals of carbon dioxide from Orbiting Carbon Observatory-2 with the version 8 ACOS algorithm, *Atmos. Meas. Tech.*, 11, 6539–6576, <https://doi.org/10.5194/amt-11-6539-2018>, 2018.
- Palmer, P. I., Feng, L., Baker, D., Chevallier, F., Bösch, H., and Somkuti, P.: Net carbon emissions from African biosphere dominate pan-tropical atmospheric CO<sub>2</sub> signal, *Nat. commun.*, 10, 1–9, <https://doi.org/10.1038/s41467-019-11097-w>, 2019.
- 1285 Peiro, H., Crowell, S., Schuh, A., Baker, D. F., O'Dell, C., Jacobson, A. R., Chevallier, F., Liu, J., Eldering, A., Crisp, D., Deng, F., Weir, B., Basu, S., Johnson, M. S., Philip, S., and Baker, I.: Four years of global carbon cycle observed from the Orbiting Carbon Observatory 2 (OCO-2) version 9 and in situ data and comparison to OCO-2 version 7, *Atmos. Chem. Phys.*, 22, 1097–1130, <https://doi.org/10.5194/acp-22-1097-2022>, 2022.
- Peters, W., van der Velde, I. R., Van Schaik, E., Miller, J. B., Ciais, P., Duarte, H. F., van der Laan-Luijckx, I. T., van der Molen, M. K.,  
1290 Scholze, M., Schaefer, K., et al.: Increased water-use efficiency and reduced CO<sub>2</sub> uptake by plants during droughts at a continental scale, *Nat. Geosci.*, 11, 744–748, <https://doi.org/10.1038/s41561-018-0212-7>, 2018.
- Petri, C., Vrekoussis, M., Rousogonous, C., Warneke, T., Sciare, J., and Notholt, J.: TCCON data from Nicosia, Cyprus (CY), Release GGG2014.R0, TCCON data archive, hosted by CaltechDATA, <https://doi.org/10.14291/tccon.ggg2014.nicosia01.R0>, 2020.
- Peylin, P., Bacour, C., MacBean, N., Leonard, S., Rayner, P., Kuppel, S., Koffi, E., Kane, A., Maignan, F., Chevallier, F., Ciais, P., and Prunet,  
1295 P.: A new stepwise carbon cycle data assimilation system using multiple data streams to constrain the simulated land surface carbon cycle, *Geosci. Model Dev.*, 9, 3321–3346, <https://doi.org/10.5194/gmd-9-3321-2016>, 2016.
- Philip, S., Johnson, M. S., Potter, C., Genovesse, V., Baker, D. F., Haynes, K. D., Henze, D. K., Liu, J., and Poulter, B.: Prior biosphere model impact on global terrestrial CO<sub>2</sub> fluxes estimated from OCO-2 retrievals, *Atmos. Chem. Phys. Discuss.*, 2019, 1–29, <https://doi.org/10.5194/acp-2018-1095>, 2019.
- 1300 Philip, S., Johnson, M. S., Baker, D. F., Basu, S., Tiwari, Y. K., Indira, N. K., Ramonet, M., and Poulter, B.: OCO-2 Satellite-Imposed Constraints on Terrestrial Biospheric CO<sub>2</sub> Fluxes Over South Asia, *J. Geophys. Res.-Atmos.*, 127, e2021JD035035, <https://doi.org/10.1029/2021JD035035>, 2022.



- Pinty, B., Janssens-Maenhout, G., Dowell, M., Zunker, H., Brunhes, T., Ciais, P., Dee, D., van der Gon, H. D., Dolman, H., Drinkwater, M., Engelen, R., Heimann, M., Holmlund, K., Husband, R., Kentarchos, A., Meijer, Y., Palmer, P., and Scholze, M.:  
1305 An operational anthropogenic CO<sub>2</sub> emissions monitoring and verification support capacity. Baseline requirements, model components and functional architecture, Tech. rep., European Commission Joint Research Centre, Brussels, <http://resolver.tudelft.nl/uuid:832e87d0-0ed8-44b2-8867-8714cebde4cb>, 2017.
- Polavarapu, S. M., Neish, M., Tanguay, M., Girard, C., Grandpré, J. d., Semeniuk, K., Gravel, S., Ren, S., Roche, S., Chan, D., et al.:  
1310 Greenhouse gas simulations with a coupled meteorological and transport model: the predictability of CO<sub>2</sub>, *Atmos. Chem. Phys.*, 16, 12 005–12 038, <https://doi.org/10.5194/acp-16-12005-2016>, 2016.
- Polavarapu, S. M., Deng, F., Byrne, B., Jones, D. B. A., and Neish, M.: A comparison of posterior atmospheric CO<sub>2</sub> adjustments obtained from in situ and GOSAT constrained flux inversions, *Atmos. Chem. Phys.*, 18, 12 011–12 044, <https://doi.org/10.5194/acp-18-12011-2018>, 2018.
- Randerson, J., Chapin, F., Harden, J., Neff, J., and Harmon, M.: Net ecosystem production: a comprehensive measure of net carbon accumulation by ecosystems, *Ecological applications*, 12, 937–947, [https://doi.org/10.1890/1051-0761\(2002\)012\[0937:NEPACM\]2.0.CO;2](https://doi.org/10.1890/1051-0761(2002)012[0937:NEPACM]2.0.CO;2), 2002.
- Rayner, P. J., Michalak, A. M., and Chevallier, F.: Fundamentals of data assimilation applied to biogeochemistry, *Atmos. Chem. Phys.*, 19, 13 911–13 932, <https://doi.org/10.5194/acp-19-13911-2019>, 2019.
- Regnier, P., Resplandy, L., Najjar, R. G., and Ciais, P.: The land-to-ocean loops of the global carbon cycle, *Nature*, 603, 401–410,  
1320 <https://doi.org/10.1038/s41586-021-04339-9>, 2022.
- Remaud, M., Chevallier, F., Cozic, A., Lin, X., and Bousquet, P.: On the impact of recent developments of the LMDz atmospheric general circulation model on the simulation of CO<sub>2</sub> transport, *Geosci. Model Dev.*, 11, 4489–4513, <https://doi.org/10.5194/gmd-11-4489-2018>, 2018.
- Remaud, M., Chevallier, F., Maignan, F., Belviso, S., Berchet, A., Parouffe, A., Abadie, C., Bacour, C., Lennartz, S., and Peylin, P.: Plant  
1325 gross primary production, plant respiration and carbonyl sulfide emissions over the globe inferred by atmospheric inverse modelling, *Atmos. Chem. Phys.*, 22, 2525–2552, <https://doi.org/10.5194/acp-22-2525-2022>, 2022.
- Ren, W., Tian, H., Tao, B., Yang, J., Pan, S., Cai, W.-J., Lohrenz, S. E., He, R., and Hopkinson, C. S.: Large increase in dissolved inorganic carbon flux from the Mississippi River to Gulf of Mexico due to climatic and anthropogenic changes over the 21st century, *J. Geophys. Res. - Biogeo.*, 120, 724–736, <https://doi.org/10.1002/2014JG002761>, 2015.
- 1330 Ren, W., Tian, H., Cai, W.-J., Lohrenz, S. E., Hopkinson, C. S., Huang, W.-J., Yang, J., Tao, B., Pan, S., and He, R.: Century-long increasing trend and variability of dissolved organic carbon export from the Mississippi River basin driven by natural and anthropogenic forcing, *Global Biogeochem. Cy.*, 30, 1288–1299, <https://doi.org/10.1002/2016GB005395>, 2016.
- Resplandy, L., Keeling, R., Rödenbeck, C., Stephens, B., Khatiwala, S., Rodgers, K., Long, M., Bopp, L., and Tans, P.: Revision of global carbon fluxes based on a reassessment of oceanic and riverine carbon transport, *Nat. Geosci.*, 11, 504–509, <https://doi.org/10.1038/s41561-018-0151-3>, 2018.
- 1335 Rödenbeck, C., Keeling, R. F., Bakker, D. C. E., Metzl, N., Olsen, A., Sabine, C., and Heimann, M.: Global surface-ocean pCO<sub>2</sub> and sea–air CO<sub>2</sub> flux variability from an observation-driven ocean mixed-layer scheme, *Ocean Sci.*, 9, 193–216, <https://doi.org/10.5194/os-9-193-2013>, 2013.
- Rödenbeck, C., Zaehle, S., Keeling, R., and Heimann, M.: How does the terrestrial carbon exchange respond to inter-annual climatic variations? A quantification based on atmospheric CO<sub>2</sub> data, *Biogeosciences*, 15, 2481–2498, <https://doi.org/10.5194/bg-15-2481-2018>, 2018.
- 1340



- Saeki, T. and Patra, P. K.: Implications of overestimated anthropogenic CO<sub>2</sub> emissions on East Asian and global land CO<sub>2</sub> flux inversion, *Geoscience Letters*, 4, 1–10, <https://doi.org/10.1186/s40562-017-0074-7>, 2017.
- Schuh, A. E., Jacobson, A. R., Basu, S., Weir, B., Baker, D., Bowman, K., Chevallier, F., Crowell, S., Davis, K. J., Deng, F., et al.: Quantifying the impact of atmospheric transport uncertainty on CO<sub>2</sub> surface flux estimates, *Global Biogeochem. Cy.*, 33, 484–500, <https://doi.org/10.1029/2018GB006086>, 2019.
- 1345 Schuh, A. E., Otte, M., Lauvaux, T., and Oda, T.: Far-field biogenic and anthropogenic emissions as a dominant source of variability in local urban carbon budgets: A global high-resolution model study with implications for satellite remote sensing, *Remote Sensing of Environment*, 262, 112473, <https://doi.org/https://doi.org/10.1016/j.rse.2021.112473>, 2021.
- Schuh, A. E., Byrne, B., Jacobson, A. R., Crowell, S. M. R., Deng, F., Baker, D. F., Johnson, M. S., Philip, S., and Weir, B.: On the role of atmospheric model transport uncertainty in estimating the Chinese land carbon sink, *Nature*, 603, E13–E14, <https://doi.org/10.1038/s41586-021-04258-9>, 2022.
- 1350 Schuld, K. N., Jacobson, A. R., Aalto, T., Andrews, A., Bakwin, P., Bergamaschi, P., Biermann, T., Biraud, S. C., Chen, H., Colomb, A., Conil, S., Cristofanelli, P., De Mazière, M., De Wekker, S., Delmotte, M., Dlugokencky, E., Emmenegger, L., Fischer, M. L., Hatakka, J., Heliasz, M., Hermanssen, O., Holst, J., Jaffe, D., Karion, A., Kazan, V., Keronen, P., Kominkova, K., Kubistin, D., Laurent, O., Laurila, T., Lee, J., Lehner, I., Leuenberger, M., Lindauer, M., Lopez, M., Mammarella, I., Manca, G., Marek, M. V., McKain, K., Miller, J. B., Miller, C. E., Myhre, C. L., Mölder, M., Müller-Williams, J., Piacentino, S., Pichon, J. M., Plass-Duelmer, C., Ramonet, M., Scheeren, B., Schumacher, M., Sha, M. K., Sloop, C. D., Smith, P., Steinbacher, M., Sweeney, C., Tans, P., Thoning, K., Trisolino, P., Tørseth, K., Viner, B., Vitkova, G., and di Sarra, A. G.: Multi-laboratory compilation of atmospheric carbon dioxide data for the years 2020–2021; obspack\_co2\_1\_NRT\_v6.1.1\_2021-05-17, <https://doi.org/10.25925/20210517>, 2021a.
- 1360 Schuld, K. N., Mund, J., Lujikx, I. T., Aalto, T., Abshire, J. B., Aikin, K., Andrews, A., Aoki, S., Apadula, F., Baier, B., Bakwin, P., Bartyzel, J., Bentz, G., Bergamaschi, P., Beyersdorf, A., Biermann, T., Biraud, S. C., Boenisch, H., Bowling, D., Brailsford, G., Chen, G., Chen, H., Chmura, L., Clark, S., Climadat, S., Colomb, A., Commane, R., Conil, S., Cox, A., Cristofanelli, P., Cuevas, E., Curcoll, R., Daube, B., Davis, K., Mazière, M. D., De Wekker, S., Della Coletta, J., Delmotte, M., DiGangi, J. P., Dlugokencky, E., Elkins, J. W., Emmenegger, L., Fang, S., Fischer, M. L., Forster, G., Frumau, A., Galkowski, M., Gatti, L. V., Gehrlein, T., Gerbig, C., Gheusi, F., Gloor, E., Gomez-Trueba, V., Goto, D., Griffis, T., Hammer, S., Hanson, C., Haszpra, L., Hatakka, J., Heimann, M., Heliasz, M., Hensen, A., Hermanssen, O., Hints, E., Holst, J., Jaffe, D., Joubert, W., Karion, A., Kawa, S. R., Kazan, V., Keeling, R., Keronen, P., Kolari, P., Kominkova, K., Kort, E., Kozlova, E., Krummel, P., Kubistin, D., Labuschagne, C., Lam, D. H., Langenfelds, R., Laurent, O., Laurila, T., Lauvaux, T., Law, B., Lee, O. S., Lee, J., Lehner, I., Leppert, R., Leuenberger, M., Levin, I., Levula, J., Lin, J., Lindauer, M., Loh, Z., Lopez, M., Machida, T., Mammarella, I., Manca, G., Manning, A., Manning, A., Marek, M. V., Martin, M. Y., Matsueda, H., McKain, K., Meijer, H., Meinhardt, F., Merchant, L., Mihalopoulos, N., Miles, N., Miller, C. E., Miller, J. B., Mitchell, L., Montzka, S., Moore, F., Morgan, E., Morgui, J.-A., Morimoto, S., Munger, B., Myhre, C. L., Mölder, M., Obersteiner, F., Müller-Williams, J., Necki, J., Newman, S., Nichol, S., Niwa, Y., O'Doherty, S., Paplawsky, B., Peischl, J., Peltola, O., Pichon, J. M., Piper, S., Plass-Duelmer, C., Ramonet, M., Ramos, R., Reyes-Sanchez, E., Richardson, S., Riris, H., Rivas, P. P., Ryerson, T., Saito, K., Sargent, M., Sasakawa, M., Sawa, Y., Say, D., Scheeren, B., Schuck, T., Schumacher, M., Seifert, T., Sha, M. K., Shepson, P., Shook, M., Sloop, C. D., Smith, P., Steinbacher, M., Stephens, B., Sweeney, C., Tans, P., Thoning, K., Timas, H., Torn, M., Trisolino, P., Turnbull, J., Tørseth, K., Vermeulen, A., Viner, B., Vitkova, G., Walker, S., Watson, A., Wofsy, S., Worsley, J., Worthy, D., Young, D., Zahn, A., Zimnoch, M., van Dinter, D., and van den Bulk, P.: Multi-laboratory compilation of atmospheric carbon dioxide data for the period 1957–2019; obspack\_co2\_1\_GLOBALVIEWplus\_v6.1\_2021-03-01, <https://doi.org/10.25925/20210301>, 2021b.
- 1370



- 1380 Sha, M. K., De Mazière, M., Notholt, J., Blumenstock, T., Chen, H., Dehn, A., Griffith, D. W. T., Hase, F., Heikkinen, P., Hermans, C., Hoffmann, A., Huebner, M., Jones, N., Kivi, R., Langerock, B., Petri, C., Scolas, F., Tu, Q., and Weidmann, D.: Intercomparison of low- and high-resolution infrared spectrometers for ground-based solar remote sensing measurements of total column concentrations of CO<sub>2</sub>, CH<sub>4</sub>, and CO, *Atmospheric Measurement Techniques*, 13, 4791–4839, <https://doi.org/10.5194/amt-13-4791-2020>, 2020.
- Sherlock, V., Connor, B. J., Robinson, J., Shiona, H., Smale, D., and Pollard, D.: TCCON data from Lauder (NZ), 120HR, Release GGG2014R0, TCCON data archive, hosted by CaltechDATA, <https://doi.org/10.14291/tcon.ggg2014.lauder01.R0/1149293>, 2014.
- 1385 Stanevich, I., Jones, D. B. A., Strong, K., Parker, R. J., Boesch, H., Wunch, D., Notholt, J., Petri, C., Warneke, T., Sussmann, R., Schneider, M., Hase, F., Kivi, R., Deutscher, N. M., Velazco, V. A., Walker, K. A., and Deng, F.: Characterizing model errors in chemical transport modeling of methane: impact of model resolution in versions v9-02 of GEOS-Chem and v35j of its adjoint model, *Geosci. Model Dev.*, 13, 3839–3862, <https://doi.org/10.5194/gmd-13-3839-2020>, 2020.
- Steinkamp, K., Mikaloff Fletcher, S. E., Brailsford, G., Smale, D., Moore, S., Keller, E. D., Baisden, W. T., Mukai, H., and Stephens, B. B.: 1390 Atmospheric CO<sub>2</sub> observations and models suggest strong carbon uptake by forests in New Zealand, *Atmos. Chem. Phys.*, 17, 47–76, <https://doi.org/10.5194/acp-17-47-2017>, 2017.
- Stephens, B. B., Gurney, K. R., Tans, P. P., Sweeney, C., Peters, W., Bruhwiler, L., Ciais, P., Ramonet, M., Bousquet, P., Nakazawa, T., et al.: Weak northern and strong tropical land carbon uptake from vertical profiles of atmospheric CO<sub>2</sub>, *Science*, 316, 1732–1735, <https://doi.org/10.1126/science.1137004>, 2007.
- 1395 Strong, K., Roche, S., Franklin, J., Mendonca, J., Lutsch, E., Weaver, D., Fogal, P., Drummond, J., Batchelor, R., and Lindenmaier, R.: TCCON data from Eureka (CA), Release GGG2014.R3 (Version R3), TCCON data archive, hosted by CaltechDATA, <https://doi.org/10.14291/TCCON.GGG2014.EUREKA01.R3>, 2019.
- Suntharalingam, P., Randerson, J. T., Krakauer, N., Logan, J. A., and Jacob, D. J.: Influence of reduced carbon emissions and oxidation on the distribution of atmospheric CO<sub>2</sub>: Implications for inversion analyses, *Global Biogeochem. Cy.*, 19, 1400 <https://doi.org/10.1029/2005GB002466>, 2005.
- Sussmann, R. and Rettinger, M.: TCCON data from Garmisch (DE), Release GGG2014.R2, <https://doi.org/10.14291/TCCON.GGG2014.GARMISCH01.R2>, 2018a.
- Sussmann, R. and Rettinger, M.: TCCON data from Zugspitze (DE), Release GGG2014R1, TCCON data archive, hosted by CaltechDATA, <https://doi.org/10.14291/tcon.ggg2014.zugspitze01.R1>, 2018b.
- 1405 Tans, P. P., Fung, I. Y., and Takahashi, T.: Observational constraints on the global atmospheric CO<sub>2</sub> budget, *Science*, 247, 1431–1438, <https://doi.org/10.1126/science.247.4949.1431>, 1990.
- Tao, B., Tian, H., Ren, W., Yang, J., Yang, Q., He, R., Cai, W., and Lohrenz, S.: Increasing Mississippi river discharge throughout the 21st century influenced by changes in climate, land use, and atmospheric CO<sub>2</sub>, *Geophys. Res. Lett.*, 41, 4978–4986, <https://doi.org/10.1002/2014GL060361>, 2014.
- 1410 Tarantola, A.: Inverse problem theory and methods for model parameter estimation, SIAM, 2005.
- Te, Y., Jeseck, P., and Janssen, C.: TCCON data from Paris (FR), Release GGG2014R0, TCCON data archive, hosted by CaltechDATA, <https://doi.org/10.14291/tcon.ggg2014.paris01.R0/1149279>, 2014.
- Thoning, K. W., Tans, P. P., and Komhyr, W. D.: Atmospheric carbon dioxide at Mauna Loa Observatory: 2. Analysis of the NOAA GMCC data, 1974–1985, *J. Geophys. Res.-Atmos.*, 94, 8549–8565, 1989.





- 1415 Tian, H., Xu, X., Liu, M., Ren, W., Zhang, C., Chen, G., and Lu, C.: Spatial and temporal patterns of CH<sub>4</sub> and N<sub>2</sub>O fluxes in terrestrial ecosystems of North America during 1979–2008: application of a global biogeochemistry model, *Biogeosciences*, 7, 2673–2694, <https://doi.org/10.5194/bg-7-2673-2010>, 2010.
- Tian, H., Ren, W., Yang, J., Tao, B., Cai, W.-J., Lohrenz, S. E., Hopkinson, C. S., Liu, M., Yang, Q., Lu, C., et al.: Climate extremes dominating seasonal and interannual variations in carbon export from the Mississippi River Basin, *Global Biogeochem. Cy.*, 29, 1333–1347, 2015a.
- 1420 Tian, H., Yang, Q., Najjar, R. G., Ren, W., Friedrichs, M. A., Hopkinson, C. S., and Pan, S.: Anthropogenic and climatic influences on carbon fluxes from eastern North America to the Atlantic Ocean: A process-based modeling study, *J. Geophys. Res. - Biogeo.*, 120, 757–772, 2015b.
- Tian, H., Xu, R., Pan, S., Yao, Y., Bian, Z., Cai, W.-J., Hopkinson, C. S., Justic, D., Lohrenz, S., Lu, C., et al.: Long-term trajectory of nitrogen loading and delivery from Mississippi River Basin to the Gulf of Mexico, *Global Biogeochem. Cy.*, 34, e2019GB006475, 2020.
- 1425 Tohjima, Y., Mukai, H., Machida, T., Nojiri, Y., and Gloor, M.: First measurements of the latitudinal atmospheric O<sub>2</sub> and CO<sub>2</sub> distributions across the western Pacific, *Geophys. Res. Lett.*, 32, 2005.
- van der Laan-Luijkx, I. T., van der Velde, I. R., van der Veen, E., Tsuruta, A., Stanislawski, K., Babenhauerheide, A., Zhang, H. F., Liu, Y., He, W., Chen, H., Masarie, K. A., Krol, M. C., and Peters, W.: The CarbonTracker Data Assimilation Shell (CTDAS) v1.0: implementation and global carbon balance 2001–2015, *Geosci. Model Dev.*, 10, 2785–2800, <https://doi.org/10.5194/gmd-10-2785-2017>, 2017.
- 1430 van der Werf, G. R., Randerson, J. T., Giglio, L., van Leeuwen, T. T., Chen, Y., Rogers, B. M., Mu, M., van Marle, M. J. E., Morton, D. C., Collatz, G. J., Yokelson, R. J., and Kasibhatla, P. S.: Global fire emissions estimates during 1997–2016, *Earth Syst. Sci. Data*, 9, 697–720, <https://doi.org/10.5194/essd-9-697-2017>, 2017.
- Walker, A. P., De Kauwe, M. G., Bastos, A., Belmecheri, S., Georgiou, K., Keeling, R. F., McMahon, S. M., Medlyn, B. E., Moore, D. J., Norby, R. J., et al.: Integrating the evidence for a terrestrial carbon sink caused by increasing atmospheric CO<sub>2</sub>, *New Phytol.*, 229, 2413–2445, 2021.
- 1435 Wang, J. A., Baccini, A., Farina, M., Randerson, J. T., and Friedl, M. A.: Disturbance suppresses the aboveground carbon sink in North American boreal forests, *Nat. Clim. Change*, 11, 435–441, 2021.
- Wang, J. S., Oda, T., Kawa, S. R., Strode, S. A., Baker, D. F., Ott, L. E., and Pawson, S.: The impacts of fossil fuel emission uncertainties and accounting for 3-D chemical CO<sub>2</sub> production on inverse natural carbon flux estimates from satellite and in situ data, *Environ. Res. Lett.*, 15, 085002, 2020.
- 1440 Warneke, T., Messerschmidt, J., Notholt, J., Weinzierl, C., Deutscher, N. M., Petri, C., and Grupe, P.: TCCON data from Orléans (FR), Release GGG2014.R1, <https://doi.org/10.14291/TCCON.GGG2014.ORLEANS01.R1>, 2019.
- Weir, B., Crisp, D., O'Dell, C. W., Basu, S., Chatterjee, A., Kolassa, J., Oda, T., Pawson, S., Poulter, B., Zhang, Z., et al.: Regional impacts of COVID-19 on carbon dioxide detected worldwide from space, *Science advances*, 7, eabf9415, 2021.
- 1445 Wennberg, P. O., Wunch, D., Roehl, C., Blavier, J.-F., Toon, G. C., and Allen, N.: TCCON data from Caltech (US), Release GGG2014R1, TCCON data archive, hosted by CaltechDATA, <https://doi.org/10.14291/tcon.ggg2014.pasadena01.R1/1182415>, 2014.
- Wennberg, P. O., Roehl, C., Blavier, J.-F., Wunch, D., Landeros, J., and Allen, N.: TCCON data from Jet Propulsion Laboratory (US), 2011, Release GGG2014R1, TCCON data archive, hosted by CaltechDATA, <https://doi.org/10.14291/tcon.ggg2014.jpl02.R1/1330096>, 2016a.
- 1450 Wennberg, P. O., Wunch, D., Roehl, C., Blavier, J.-F., Toon, G. C., Allen, N., Dowell, P., Teske, K., Martin, C., and Martin, J.: TCCON data from Lamont (US), Release GGG2014R1, TCCON data archive, hosted by CaltechDATA, <https://doi.org/10.14291/tcon.ggg2014.lamont01.R1/1255070>, 2016b.



- Wennberg, P. O., Roehl, C. M., Wunch, D., Toon, G. C., Blavier, J.-F., Washenfelder, R., Keppel-Aleks, G., Allen, N. T., and Ayers, J.: TCCON data from Park Falls (US), Release GGG2014.R1, <https://doi.org/10.14291/TCCON.GGG2014.PARKFALLS01.R1>, 2017.
- 1455 Worden, J. R., Cusworth, D. H., Qu, Z., Yin, Y., Zhang, Y., Bloom, A. A., Ma, S., Byrne, B. K., Scarpelli, T., Maasackers, J. D., Crisp, D., Duren, R., and Jacob, D. J.: The 2019 methane budget and uncertainties at 1° resolution and each country through Bayesian integration Of GOSAT total column methane data and a priori inventory estimates, *Atmos. Chem. Phys.*, 22, 6811–6841, <https://doi.org/10.5194/acp-22-6811-2022>, 2022.
- 1460 Wunch, D., Toon, G. C., Wennberg, P. O., Wofsy, S. C., Stephens, B. B., Fischer, M. L., Uchino, O., Abshire, J. B., Bernath, P., Biraud, S. C., Blavier, J.-F. L., Boone, C., Bowman, K. P., Browell, E. V., Campos, T., Connor, B. J., Daube, B. C., Deutscher, N. M., Diao, M., Elkins, J. W., Gerbig, C., Gottlieb, E., Griffith, D. W. T., Hurst, D. F., Jiménez, R., Keppel-Aleks, G., Kort, E. A., Macatangay, R., Machida, T., Matsueda, H., Moore, F., Morino, I., Park, S., Robinson, J., Roehl, C. M., Sawa, Y., Sherlock, V., Sweeney, C., Tanaka, T., and Zondlo, M. A.: Calibration of the Total Carbon Column Observing Network using aircraft profile data, *Atmos. Meas. Tech.*, 3, 1351–1362, <https://doi.org/10.5194/amt-3-1351-2010>, 2010.
- 1465 Wunch, D., Toon, G. C., Blavier, J.-F. L., Washenfelder, R. A., Notholt, J., Connor, B. J., Griffith, D. W., Sherlock, V., and Wennberg, P. O.: The Total Carbon Column Observing Network, *Philos. T. Roy. Soc. A*, 369, 2087–2112, <https://doi.org/10.1098/rsta.2010.0240>, 2011.
- Wunch, D., Mendonca, J., Colebatch, O., Allen, N., Blavier, J.-F. L., Roche, S., Hedelius, J. K., Neufeld, G., Springett, S., Worthy, D. E. J., Kessler, R., and Strong, K.: TCCON data from East Trout Lake (CA), Release GGG2014R1, TCCON data archive, hosted by CaltechDATA, <https://doi.org/10.14291/tcon.ggg2014.easttroutlake01.R1>, 2017a.
- 1470 Wunch, D., Wennberg, P. O., Osterman, G., Fisher, B., Naylor, B., Roehl, C. M., O'Dell, C., Mandrake, L., Viatte, C., Kiel, M., Griffith, D. W. T., Deutscher, N. M., Velasco, V. A., Notholt, J., Warneke, T., Petri, C., De Maziere, M., Sha, M. K., Sussmann, R., Rettinger, M., Pollard, D., Robinson, J., Morino, I., Uchino, O., Hase, F., Blumenstock, T., Feist, D. G., Arnold, S. G., Strong, K., Mendonca, J., Kivi, R., Heikkinen, P., Iraci, L., Podolske, J., Hillyard, P. W., Kawakami, S., Dubey, M. K., Parker, H. A., Sepulveda, E., García, O. E., Te, Y., Jeseck, P., Gunson, M. R., Crisp, D., and Eldering, A.: Comparisons of the Orbiting Carbon Observatory-2 (OCO-2)  $X_{CO_2}$  measurements with TCCON, *Atmos. Meas. Tech.*, 10, 2209–2238, <https://doi.org/10.5194/amt-10-2209-2017>, 2017b.
- 1475 Yang, D., Liu, Y., Cai, Z., Wang, M., Qiu, L., Yin, Z., Tian, L., and TanSat-2 team: The next generation of Chinese greenhouse gas monitoring satellite mission, 14th International Workshop on Greenhouse Gas Measurements from Space (IWGGMS-14), 2018.
- Yang, Q., Tian, H., Friedrichs, M. A., Hopkinson, C. S., Lu, C., and Najjar, R. G.: Increased nitrogen export from eastern North America to the Atlantic Ocean due to climatic and anthropogenic changes during 1901–2008, *J. Geophys. Res. - Biogeo.*, 120, 1046–1068, 2015.
- 1480 Yao, Y., Tian, H., Shi, H., Pan, S., Xu, R., Pan, N., and Canadell, J. G.: Increased global nitrous oxide emissions from streams and rivers in the Anthropocene, *Nat. Clim. Change*, 10, 138–142, 2020.
- Yao, Y., Tian, H., Pan, S., Najjar, R. G., Friedrichs, M. A., Bian, Z., Li, H.-Y., and Hofmann, E. E.: Riverine carbon cycling over the past century in the Mid-Atlantic region of the United States, *J. Geophys. Res. - Biogeo.*, 126, e2020JG005968, 2021.
- 1485 Yin, Y., Ciais, P., Chevallier, F., Van der Werf, G. R., Fanin, T., Broquet, G., Boesch, H., Cozic, A., Hauglustaine, D., Szopa, S., et al.: Variability of fire carbon emissions in equatorial Asia and its nonlinear sensitivity to El Niño, *Geophys. Res. Lett.*, 43, 10–472, 2016.
- Yu, K., Keller, C. A., Jacob, D. J., Molod, A. M., Eastham, S. D., and Long, M. S.: Errors and improvements in the use of archived meteorological data for chemical transport modeling: an analysis using GEOS-Chem v11-01 driven by GEOS-5 meteorology, *Geosci. Model Dev.*, 11, 305–319, <https://doi.org/10.5194/gmd-11-305-2018>, 2018.
- 1490 Zammit-Mangion, A., Bertolacci, M., Fisher, J., Stavert, A., Rigby, M., Cao, Y., and Cressie, N.: WOMBAT v1.0: A Fully Bayesian Global Flux-Inversion Framework, *Geosci. Model Dev.*, 15, 45–73, <https://doi.org/10.5194/gmd-15-45-2022>, 2022.



Zeng, J., Nojiri, Y., Landschützer, P., Telszewski, M., and Nakaoka, S.-I.: A global surface ocean  $f\text{CO}_2$  climatology based on a feed-forward neural network, *J. Atmos. Ocean Tech.*, 31, 1838–1849, 2014.

Zhang, L., Davis, K. J., Schuh, A. E., Jacobson, A. R., Pal, S., Cui, Y. Y., Baker, D., Crowell, S., Chevallier, F., Remaud, M., et al.: Multi-Season Evaluation of  $\text{CO}_2$  Weather in OCO-2 MIP Models, *J. Geophys. Res.-Atmos.*, 127, e2021JD035 457, 2022.