

Copernicus Arctic Regional Reanalysis (CARRA): list of references

Last modified on Apr 30, 2024 12:38

Table of Contents

- Peer-reviewed studies
- A selection of relevant presentations
- A selection of relevant conference abstracts and not peer-reviewed literature
- Relevant CARRA documents
- Related articles

This page includes a list of scientific work that fully or partly include diagnostics and verification and/or description of the Copernicus Arctic Regional Reanalysis (CARRA) data set. The list includes links, abstract of peer-reviewed work and some details to provide initial insight on what part of the CARRA data is evaluated. The list is not necessarily complete in terms of including all published work on and with the CARRA data set, but provides a starting point to seek information on the quality of different aspects of the data set in the literature.

Peer-reviewed studies

Schyberg et al. in preparation: The Copernicus Arctic regional reanalysis				
Type of study: System description	Parameters: 2m air temperature and humidity, 10m wind speed, precipitation, Mean sea Level Pressure	Comparison against: synop observations, ERA5	Region and time period: 1991-2020, both CARRA domains (sub-regions: Svalbard, coast, inland etc...)	Features: Summary statistics, trends, case-studies
Summary/abstract (paper): This paper is in preparation and will provide a full scientific description and evaluation of the Copernicus Arctic Regional Reanalysis. Abstract and link to published paper will be provided when published.				

Batrak, Y., Cheng, B., and Kallio-Myers, V.: Sea ice cover in the Copernicus Arctic Regional Reanalysis, The Cryosphere Discuss. [preprint], https://doi.org/10.5194/tc-2023-74 , in review, 2023.				
Type of study: Evaluation /verification	Parameters: Sea ice properties	Comparison against: satellite products, buoys, ERA5	Region and time period: mainly 2000-2020, CARRA-East and CARRA-West	Features: Summary statistics, climatology
Summary/abstract (paper): The Copernicus Arctic Regional Reanalysis (CARRA) is a novel regional high-resolution atmospheric reanalysis product that covers a considerable part of the European Arctic including substantial amounts of ice-covered areas. Sea ice in CARRA is modelled by means of a one-dimensional thermodynamic sea ice parameterisation scheme, which also explicitly resolves the evolution of the snow layer over sea ice. In the present study we assess the representation of sea ice cover in the CARRA product and validate it against a wide set of satellite products and observations from ice mass balance buoys. We show that sea ice cover in CARRA adequately represents general interannual trends towards thinner and warmer ice in the Arctic. Compared to ERA5, sea ice in CARRA shows a reduced warm bias in the ice surface temperature. The strongest improvement was observed for winter months over the Central Arctic, and the Greenland and Barents seas where a 4.91 °C median ice surface temperature error of ERA5 is reduced to 1.88 °C in CARRA on average. Over the Baffin Bay, intercomparisons suggest the presence of a cold winter-time ice surface temperature bias in CARRA. No improvement over ERA5 was found in the ice surface albedo with spring-time errors in CARRA being up to 8 % higher on average than those in ERA5 when computed against the CLARA-A2 satellite retrieval product. Summer-time ice surface albedos are comparable in CARRA and ERA5. Sea ice thickness and snow depth in CARRA adequately resolve the annual cycle of sea ice cover in the Arctic and bring added value compared to ERA5. However, limitations of CARRA indicate potential benefits of utilising more advanced approaches for representing sea ice cover in next generation reanalyses. https://tc.copernicus.org/preprints/tc-2023-74/				

Box, J. E., Nielsen, K. P., Yang, X., Niwano, M., Wehrli, A., van As, D., Fettweis, X., Køltzow, Morten A. Ø., Palmason, B., Fausto, R. S., van den Broeke, M. R., Huai, B., Ahlstrøm, A. P., Langley, K., Dachauer, A., & Noël, B. (2023). Greenland ice sheet rainfall climatology, extremes and atmospheric river rapids. <i>Meteorological Applications</i> , 30(4), e2134. https://doi.org/10.1002/met.2134				
Type of study: Evaluation and precipitation analysis	Parameters: Rainfall	Comparison against: in-situ, ERA5	Region and time period: CARRA -West	Features: Summary statistics, climatology, trends, case studies
Summary/abstract (paper): Greenland rainfall has come into focus as a climate change indicator and from a variety of emerging cryospheric impacts. This study first evaluates rainfall in five state-of-the-art numerical prediction systems (NPSs) (CARRA, ERA5, NHM-SMAP, RACMO, MAR) using in situ rainfall data from two regions spanning from land onto the ice sheet. The new EU Copernicus Climate Change Service (C3S) Arctic Regional ReAnalysis (CARRA), with a relatively fine (2.5km) horizontal grid spacing and extensive within-model-domain observational initialization, has the lowest average bias and highest explained variance relative to the field data. ERA5 inland wet bias versus CARRA is consistent with the field data and other research and is presumably due to more ERA5 topographic smoothing. A CARRA climatology 1991–2021 has rainfall increasing by more than one-third for the ice sheet and its peripheral ice masses. CARRA and in situ data illuminate extreme (above 300mm per day) local rainfall episodes. A detailed examination CARRA data reveals the interplay of mass conservation that splits flow around southern Greenland and condensational buoyancy generation that maintains along-flow updraft ‘rapids’ 2km above sea level, which produce rain bands within an atmospheric river interacting with Greenland. CARRA resolves gravity wave oscillations that initiate as a result of buoyancy offshore, which then amplify from terrain-forced uplift. In a detailed case study, CARRA resolves orographic intensification of rainfall by up to a factor of four, which is consistent with the field data. https://rmets.onlinelibrary.wiley.com/doi/full/10.1002/met.2134				

Frank, L., Jonassen, M. O., Skogseth, R., & Vihma, T. (2023). Atmospheric climatologies over Isfjorden, Svalbard. <i>Journal of Geophysical Research: Atmospheres</i> , 128, e2022JD038011. https://doi.org/10.1029/2022JD038011				
---	--	--	--	--

Type of study: climatology	Parameters: near-surface temperature, humidity and wind as well as total cloud cover and precipitation	Comparison against:	Region and time period: Isfjorden, Svalbard, June 2011–May 2021	Features: climate statistics and synoptic flow type
Summary/abstract (paper): <p>The Isfjorden region at the west coast of Spitsbergen is the most easily accessible area in the Svalbard Archipelago, making it a perfect outdoor laboratory for Arctic research. Due to its location in the high Arctic together with its complex terrain, the climatic conditions vary substantially both in time and space. Based on a new high-resolution reanalysis, we present climatologies of five major atmospheric variables over the Isfjorden region during 2011–2021 with special focus on local effects. For example, we find that topographic channeling effects often lead to differences in near-surface wind speeds of several m s⁻¹ over small horizontal distances. During winter, the fjord acts as a heat and moisture island, ultimately impacting the adjacent low-elevation land areas. These land–sea gradients reverse during summer. High mountain areas surrounding the fjord experience substantially different climatic conditions, with for example, seasonal precipitation doubling from sea level to approximately 700 m. The spatial variability over the Isfjorden region is in general found to be smaller than its temporal counterpart but larger than the diurnal cycle. Besides these findings, this study furthermore demonstrates the importance of high-resolution regional atmospheric reanalyses compared to global products for the characterization of the local micro-climate over Arctic fjords and the interaction with surrounding land areas.</p> <p>https://doi.org/10.1029/2022JD038011</p>				
Grinsted, A., Rathmann, N. M., Mottram, R., Solgaard, A. M., Mathiesen, J., and Hvidberg, C. S.: Failure strength of glacier ice inferred from Greenland crevasses, EGUsphere [preprint], https://doi.org/10.5194/egusphere-2023-1957, 2023				
Type of study: Glacier	Parameters: surface air temperature	Comparison against:	Region and time period: Greenland, 1991–2020	Features: Input to glacier calculations
Summary/abstract (paper): Ice fractures when subject to stress that exceeds the material failure strength. Previous studies have found that a von Mises failure criterion, which places a bound on the second invariant of the deviatoric stress tensor, is consistent with empirical data. Other studies have suggested that a scaling effect exists, such that larger sample specimens have a substantially lower failure strength, implying that estimating material strength from laboratory-scale experiments may be insufficient for glacier-scale modelling. In this paper, we analyze the stress conditions in crevasse onset regions to better understand the failure criterion and strength relevant for large-scale modelling. The local deviatoric stress is inferred using surface velocities and reanalysis temperatures, and crevasse onset regions are extracted from a remotely sensed crevasse density map. We project the stress state onto the failure plane spanned by Haigh–Westergaard coordinates, showing how failure depends on mode of stress. We find that existing crevasse data is consistent with a Schmidt–Ishlinsky failure criterion that places a bound on the absolute value of the maximal principal deviatoric stress, estimated to be (158 ± 44) kPa. Although the traditional von Mises failure criterion also provides an adequate fit to the data with a von Mises strength of (265 ± 73) kPa, it depends only on stress magnitude and is indifferent to the specific stress state, unlike Schmidt–Ishlinsky failure which has a larger shear failure strength compared to tensile strength. Implications for large-scale ice-flow and fracture modelling are discussed. <p>https://doi.org/10.5194/egusphere-2023-1957</p>				
Hansche, I., Shahi, S., Abermann, J., & Schöner, W. (2023). The vertical atmospheric structure of the partially glacierised Mittivakkat valley, southeast Greenland. Journal of Glaciology, 1-12. doi:10.1017/jog.2022.120				
Type of study: Investigation of atmospheric temperature inversions	Parameters: air temperature	Comparison against: UAV-observations compared with CARRA, ERA5, ERA-interim and radiosondes	Region and time period: Mittivakkat valley (southeast Greenland), field campaign summer 2019	Features; Inversion characteristics, comparisons by correlation and RMSE
Summary/abstract(paper): <p>Air temperature inversions, a situation in which atmospheric temperature increases with height, are key components of the Arctic planetary boundary layer. The present study investigates the spatial and temporal variations of temperature inversions over different surface types (rock, gravel, snow, ice) along the Mittivakkat valley (southeast Greenland). For this purpose, 113 vertical profiles with high spatio-temporal resolution of air temperature and relative humidity were collected with unoccupied aerial vehicles (UAVs) during a 13-day field campaign in summer 2019. Air temperature inversions were present in 83% of the profiles, of which 24% were surface-based inversions and 76% were elevated inversions. The proglacial area covered with bare rock and gravel induces surface heating and convection during the day and, through interaction with local circulation patterns, leads to the frequent formation of elevated inversions. In contrast, the glacier surface itself acts as a persistent cooling surface and leads to the formation of surface-based inversions. A low-level fog layer that forms under the inversion layer may be causing non-linear vertical ablation gradients on Mittivakkat Gletsjer. Furthermore, we demonstrate that atmospheric measurements using UAVs can better capture small-scale processes than other products like radiosonde or modeled reanalysis data.</p> <p>https://www.cambridge.org/core/journals/journal-of-glaciology/article/vertical-atmospheric-structure-of-the-partially-glacierised-mittivakkat-valley-southeast-greenland/F47255DD27CFAE332D23FD61FBEEB448</p>				
Helgason, H. B. and Nijssen, B.: LamaH-Ice: LARge-SaMple Data for Hydrology and Environmental Sciences for Iceland, Earth Syst. Sci. Data Discuss. [preprint], https://doi.org/10.5194/essd-2023-349, in review, 2023				
Type of study: Hydrology	Parameters: Precipitation	Comparison against: in-situ observations	Region and time period: Iceland, 1991–2023	Features: Precipitation amount
Summary/abstract (paper): <p>Access to mountainous regions for monitoring streamflow, snow and glaciers is often difficult, and many rivers are thus not gauged and hydrological measurements are limited. Consequently, cold-region watersheds, particularly heavily glacierized ones, are poorly represented in large-sample hydrology (LSH) datasets. We present a new LSH dataset for Iceland, termed LamaH-Ice (LARge-SaMple DAta for Hydrology and Environmental Sciences for Iceland). Glaciers and ice caps cover about 10 % of Iceland and while streamflow has been measured for several decades, these measurements have not been published in a consistent manner before. The dataset provides daily and hourly hydrometeorological timeseries and catchment characteristics for 107 river basins in Iceland, covering an area of almost 46,000 km² (45 % of Iceland's area), with catchment sizes ranging from 4 to about 7,500 km². LamaH-Ice conforms to the structure of existing LSH datasets and includes most variables offered in these datasets, as well as additional information relevant to cold-region hydrology, e.g., timeseries of snow cover, glacier mass balance and albedo. LamaH-Ice also includes dynamic catchment characteristics to account for changes in land cover, vegetation, and glacier extent. A large majority of the watersheds in LamaH-Ice are not subject to human activities, such as diversions and flow regulations. Streamflow measurements under natural flow conditions are highly valuable to hydrologists seeking to model and comprehend the natural hydrological cycle or estimate climate change trends. The LamaH-Ice dataset (Helgason and Nijssen, 2023) is intended for the research community to improve the understanding of hydrology in cold-region environments.</p> <p>https://doi.org/10.5194/essd-2023-349</p>				
Khachatryan, E.; Asemann, P.; Zhou, L.; Birkelund, Y.; Esau, I.; Ricaud, B. Exploring the Potential of Sentinel-1 Ocean Wind Field Product for Near-Surface Offshore Wind Assessment in the Norwegian Arctic. Atmosphere 2024, 15, 146. https://doi.org/10.3390/atmos15020146				
Type of study: Intercomparison	Parameters: wind speed	Comparison against: Sentinel wind, in-situ, ERA5, NORA3	Region and time period: Goliat station off the Northern Norway coast, 2022	Features: wind

Summary/abstract (paper):

The exploitation of offshore wind resources is a crucial step towards a clean energy future. It requires an advanced approach for high-resolution wind resource evaluations. We explored the suitability of the Sentinel-1 Level-2 OCN ocean wind field (OWI) product for offshore wind resource assessments. The SAR data were compared to in situ observations and three reanalysis products: the global reanalysis ERA5 and two regional reanalyses CARRA and NORA3. This case study matches 238 scenes from 2022 for the Goliat station, an oil platform located 85 km northwest of Hammerfest in the Barents Sea, where a new offshore wind park has been proposed. The analysis showed that despite their unique limitations in spatial and temporal resolutions, all data sources have similar statistical properties (RMSE, correlation coefficient, and standard deviation). The Weibull parameters characterizing the wind speed distributions showed strong similarities between the Sentinel-1 and all reanalysis data. The Weibull parameters of the in situ measurements showed an underestimation of wind speed compared to all other sources. Comparing the full reanalysis datasets with the subsets matching the SAR scenes, only slight changes in Weibull parameters were found, indicating that, despite its low temporal resolution, the Sentinel-1 Level 2 OWI product can compete with the more commonly used reanalysis products in the estimation of offshore wind resources. Its high spatial resolution, which is unmatched by other methods, renders it especially valuable in offshore areas close to complex coastlines and in resolving weather events at a smaller scale.

<https://doi.org/10.3390/atmos15020146>



Kirbus, B., Schirmacher, I., Klingebiel, M., Schäfer, M., Ehrlich, A., Slättberg, N., Lucke, J., Moser, M., Müller, H., and Wendisch, M.: Thermodynamic and cloud evolution in a cold air outbreak during HALO-(AC)3: Quasi-Lagrangian observations compared to the ERA5 and CARRA reanalyses, EGUsphere [preprint], <https://doi.org/10.5194/egusphere-2023-2989>, 2023.

Type of study: Arctic air mass transformations	Parameters: temperature, humidity, clouds	Comparison against: in-situ observations and ERA5	Region and time period: Fram strait, 1. April 2022	Features: Cold air outbreak
---	--	--	---	------------------------------------

Summary/abstract (paper):

Intense air mass transformations take place when cold, dry Arctic air masses move southward from the closed sea ice onto the much warmer ice-free Arctic ocean during marine cold air outbreaks (MCAOs). In spite of intensive research on MCAOs during recent years, the temporal rates of diabatic heating and moisture uptake relevant also for cloud formation/dissipation have not been measured along MCAO flows. Instead, reanalyses have typically been used for climatological investigations of MCAOs or to supply higher-resolution models with lateral boundary conditions and time-dependent forcings. Meanwhile, the uncertainties connected to those datasets remain unclear.

Here, we present height-resolved observations of diabatic heating rates, moisture uptake, and cloud evolution measured in a quasi-Lagrangian manner. The investigated specific MCAO was observed on 01 April 2022 during the HALO-(AC)3 airborne campaign that was conducted in spring 2022. Shortly after passing the ice edge, maximum diabatic heating rates larger than 6 K h⁻¹ and moisture uptake of more than 0.3 g kg⁻¹ h⁻¹ were measured close above the ocean surface. As the air mass continued its drift southwards, clouds started to form and vertical mixing within the steadily deepening boundary layer was intensified. The quasi-Lagrange observations are compared with reanalysis data from the European Centre for Medium-Range Weather Forecasts (ECMWF) latest global reanalysis ERA5 and the Copernicus Arctic Regional Reanalysis (CARRA). It was found that the mean absolute errors (MAEs) of ERA5 versus CARRA data are 60 % higher for air temperature over sea ice (1.4 K versus 0.9 K), and 70 % higher for specific humidity over ice-free ocean (0.12 g kg⁻¹ versus 0.07 g kg⁻¹). We relate these differences not only to issues with representations of the marginal ice zone and corresponding surface fluxes in ERA5, but also to the cloud scheme producing excess liquid-bearing clouds and precipitation, causing a too-dry marine boundary layer. Overall, the combination of CARRA's high spatial resolution, an improved handling of cold surfaces, and the demonstrated higher fidelity towards the observations, make it a well-suited candidate for further investigations of Arctic air mass transformations.

<https://doi.org/10.5194/egusphere-2023-2989>

Køltzow M., Schyberg H., Støylen E., & Yang X. (2022). Value of the Copernicus Arctic Regional Reanalysis (CARRA) in representing near-surface temperature and wind speed in the north-east European Arctic. Polar Research, 41. <https://doi.org/10.33265/polar.v41.8002>

Type of study: Eval uation/verification	Parameters: 2m air temperature, 10m wind speed	Comparison against: syno p observations, ERA5	Region and time period: 1998-2018, CARRA-East (s ub-regions: Svalbard, coast, inland etc...)	Features: Summary statistics, spatial and temporal variability, extremes/ high-impact
--	---	--	---	--

Summary/abstract (paper):

The representation of 2-m air temperature and 10-m wind speed in the high-resolution (with a 2.5-km grid spacing) Copernicus Arctic Regional Reanalysis (CARRA) and the coarser resolution (ca. 31-km grid spacing) global European Center for Medium-range Weather Forecasts fifth-generation reanalysis (ERA5) for Svalbard, northern Norway, Sweden and Finland is evaluated against observations. The largest differences between the two reanalyses are found in regions with complex terrain and coastlines, and over the sea ice for temperature in winter. In most aspects, CARRA outperforms ERA5 in its agreement with the observations, but the value added by CARRA varies with region and season. Furthermore, the added value by CARRA is seen for both parameters but is more pronounced for temperature than wind speed. CARRA is in better agreement with observations in terms of general evaluation metrics like bias and standard deviation of the errors, is more similar to the observed spatial and temporal variability and better captures local extremes. A better representation of high-impact weather like polar lows, vessel icing and warm spells during winter is also demonstrated. Finally, it is shown that a substantial part of the difference between reanalyses and observations is due to representativeness issues, that is, sub-grid variability, which cannot be represented in gridded data. This representativeness error is larger in ERA5 than in CARRA, but the fraction of the total error is estimated to be similar in the two analyses for temperature but larger in ERA5 for wind speed.

<https://polarresearch.net/index.php/polar/article/view/8002/14479>

Isaksen, K., Nordli, Ø., Ivanov, B. et al. Exceptional warming over the Barents area. Sci Rep 12, 9371 (2022). <https://doi.org/10.1038/s41598-022-13568-5>

Type of study:	Parameters:	Comparison against:	Region and time period:	Features;
Investigation of temperature trends	2m air temperature	ERA5, dependent and in-dependent in-situ/synop	1991-2020, Svalbard and Barents Sea area	Trends, summary statistics (bias, standard deviation of error, RMSE), sea-ice dependency

Summary/abstract(paper):

In recent decades, surface air temperature (SAT) data from Global reanalyses points to maximum warming over the northern Barents area. However, a scarcity of observations hampers the confidence of reanalyses in this Arctic hotspot region. Here, we study the warming over the past 20–40 years based on new available SAT observations and a quality controlled comprehensive SAT dataset from the northern archipelagos in the Barents Sea. We identify a statistically significant record-high annual warming of up to 2.7 °C per decade, with a maximum in autumn of up to 4.0 °C per decade. Our results are compared with the most recent global and Arctic regional reanalysis data sets, as well as remote sensing data records of sea ice concentration (SIC), sea surface temperature (SST) and high-resolution ice charts. The warming pattern is primarily consistent with reductions in sea ice cover and confirms the general spatial and temporal patterns represented by reanalyses. However, our findings suggest even a stronger rate of warming and SIC-SAT relation than was known in this region until now.

<https://www.nature.com/articles/s41598-022-13568-5>

Moore, G. W. K., & Imrit, A. A. (2022). Impact of resolution on the representation of the mean and extreme winds along Nares Strait. Journal of Geophysical Research: Atmospheres, 127, e2022JD037443. <https://doi.org/10.1029/2022JD037443>

Type of study:	Parameters:	Comparison against:	Region and time period:	Features;
Investigation of wind speed in complex terrain	10 m wind speed	ERA5, ECMWF Operational analysis and in-situ/synop	2016-2019, Nares strait	General statistics and extremes

Multiple authors, 2020: [User workshop on Copernicus regional reanalysis for Europe and the European Arctic, September 2020](#). Multiple (recorded) presentations.

A selection of relevant conference abstracts and not peer-reviewed literature

Dahlgren, P. and T. Valkonen, 2021: Use of wind retrievals in regional reanalysis, 15th International Winds Workshop, online, abstract available in abstract brochure here: <http://cimss.ssec.wisc.edu/iwwg/iww15/index.html>

Kallio-Myers, V., Batrak, Y., and Cheng, B.: Comparison of Arctic sea-ice albedo between CARRA and ERA5 reanalyses and satellite based CLARA-A2, EGU General Assembly 2023, Vienna, Austria, 24–28 Apr 2023, EGU23-1510, <https://doi.org/10.5194/egusphere-egu23-1510>, 2023.

Landgren, O., Lutz, J., Dobler, A., and Isaksen, K.: Multi-decadal convection-permitting climate simulation over Svalbard and its benefit for assessing the future of cultural heritage sites, EMS Annual Meeting 2022, Bonn, Germany, 5–9 Sep 2022, EMS2022-556, <https://doi.org/10.5194/ems2022-556>, 2022

Maniktala, D., 2022, Analysing seasonal snow cover trends and patterns on Svalbard, student thesis, Uppsala University, Disciplinary Domain of Science and Technology, Earth Sciences, Department of Earth Sciences, LUVAL. <https://www.diva-portal.org/smash/record.jsf?pid=diva2:1689663>

Nielsen, K. P., Schyberg, H., Yang, X., Støylen, E., Dahlgren, P., Amstrup, B., Peralta, C., Køltzow, M., and Bojarova, J.: 24 years of C3S Arctic regional reanalysis, EGU General Assembly 2021, online, 19–30 Apr 2021, EGU21-15178, <https://doi.org/10.5194/egusphere-egu21-15178>, 2021.

Schyberg, H. The Copernicus Arctic Regional Reanalysis, WCRP-WWRP Symposium on Data Assimilation and Reanalysis / 2021 ECMWF Annual Seminar on Observations, 13-18 September 202, <https://symp-bonn2021.sciencesconf.org/data/357176.pdf>

Schyberg, H., Yang, X., Støylen, E., Dahlgren, P., S. Madsen, M., Køltzow, M., and Olesen, M.: Evolution of the Copernicus Arctic Regional Reanalysis, EMS Annual Meeting 2022, Bonn, Germany, 5–9 Sep 2022, EMS2022-535, <https://doi.org/10.5194/ems2022-535>, 2022.

Slättberg, N., Maturilli, M., and Dahlke, S.: Fram Strait Marine Cold Air Outbreaks and associated surface heat fluxes in the ERA5 & CARRA reanalyses, EGU General Assembly 2023, Vienna, Austria, 24–28 Apr 2023, EGU23-14048, <https://doi.org/10.5194/egusphere-egu23-14048>, 2023.

Torres-Alavez, A., Landgren, O., Boberg, F., Christensen, O. B., Mottram, R., Olesen, M., Van Ulft, B., Verro, K., and Batrak, Y.: Assessing Performance of a new High Resolution polar regional climate model with remote sensing and in-situ observations: HCLIM in the Arctic and Antarctica, EGU General Assembly 2023, Vienna, Austria, 24–28 Apr 2023, EGU23-14090, <https://doi.org/10.5194/egusphere-egu23-14090>, 2023.

Zhaohui, Cheng: Polar mesoscale cyclones in ERA5 and CARRA, 2023, Uppsala University, Disciplinary Domain of Science and Technology, Earth Sciences, Department of Earth Sciences, LUVAL. <https://www.diva-portal.org/smash/record.jsf?pid=diva2%3A1765122&dswid=6826>

Relevant CARRA documents

Nielsen, K. P. et al.: Copernicus Arctic Regional Reanalysis (CARRA): Data User Guide. Available at [Copernicus Arctic Regional Reanalysis \(CARRA\): Data User Guide](#)

Yang, X., et al., 2020: C3S Arctic regional reanalysis - Full System documentation. Available at <https://datastore.copernicus-climate.eu/documents/reanalysis-carra/CARRAFullSystemDocumentationFinal.pdf>

Yang, X., et al., 2020: Complete test and verification report on fully configured reanalysis and monitoring system. Available at <https://datastore.copernicus-climate.eu/documents/reanalysis-carra/CARRATestVerificationFinal.pdf>

Bojarova J., 2020: Uncertainty estimation method. Available at <https://datastore.copernicus-climate.eu/documents/reanalysis-carra/CARRAUncertainty%20estimationFinal.pdf>

Copernicus Arctic Regional Reanalysis: Added value to the ERA5 global reanalysis. Copernicus Knowledge Base (CKB) article. [Copernicus Arctic Regional Reanalysis \(CARRA\): Added value to the ERA5 global reanalysis](#)

Uncertainty information for the Copernicus Arctic Regional reanalysis. Copernicus Knowledge Base (CKB) article. [Copernicus Arctic Regional Reanalysis \(CARRA\): known issues and uncertainty information#Uncertaintyinformation](#)

Known issues for the Copernicus Arctic Regional reanalysis. Copernicus Knowledge Base (CKB) article. [Copernicus Arctic Regional Reanalysis \(CARRA\): known issues and uncertainty information#Knownissues](#)

This document has been produced in the context of the Copernicus Climate Change Service (C3S).

The activities leading to these results have been contracted by the European Centre for Medium-Range Weather Forecasts, operator of C3S on behalf of the European Union (Delegation agreement signed on 11/11/2014). All information in this document is provided "as is" and no guarantee or warranty is given that the information is fit for any particular purpose.

The users thereof use the information at their sole risk and liability. For the avoidance of all doubt, the European Commission and the European Centre for Medium - Range Weather Forecasts have no liability in respect of this document, which is merely representing the author's view.

Related articles

- [Copernicus Arctic Regional Reanalysis \(CARRA\): known issues and uncertainty information](#)

- [Copernicus Arctic Regional Reanalysis \(CARRA\): list of references](#)
- [Copernicus Arctic Regional Reanalysis \(CARRA\): Added value to the ERA5 global reanalysis](#)
- [Copernicus Arctic Regional Reanalysis \(CARRA\): Data User Guide](#)
- [Copernicus Arctic Regional Reanalysis \(CARRA\)](#)